

A checklist of species of Cylindroleberididae (Crustacea: Ostracoda)

ANNA E. SYME AND GARY C. B. POORE

Museum Victoria, GPO Box 666, Melbourne, Victoria, 3001, Australia
(asyme@museum.vic.gov.au)

Abstract

Syme, A.E., and Poore, G.C.B. 2006. A checklist of species of Cylindroleberididae (Crustacea: Ostracoda). *Museum Victoria Science Reports* 9: 1–20.

Presented here is a checklist for the Cylindroleberididae, including a list of current genera, species, synonyms, type localities, and remarks concerning taxonomic uncertainties. A taxonomic history of the family is provided, along with a listing of higher systematics. There are currently three subfamilies, five tribes, 32 genera and 219 described species. A checklist is a necessary starting point for a future systematic revision of the Cylindroleberididae.

Keywords

Crustacea, Ostracoda, Cylindroleberididae, checklist

Introduction

Cylindroleberidid ostracods are found in the marine environment at all latitudes and depths. Currently there are 219 described species in 32 genera. As a family they are well-defined by several morphological features, including gills. Early monographs described the family and known species in detail (Skogsberg, 1920; Poulsen, 1965). Major monographs of the last 50 years are by Kornicker (e.g., 1975, 1981b, 1986). For all ostracods, a large database in several volumes is available by Kempf, (1986-2004), which lists references to all described genera and species. However, this database can only be searched by species or genera; it is not searchable by higher taxonomic categories such as family.

This report provides a reference point for those wishing to identify cylindroleberidid ostracods and is a basis for a future systematic revision. It includes a checklist of genera and species of Cylindroleberididae, including type species and synonyms. Although the family is well-defined, there is some uncertainty regarding the types of some genera and species. For example, the holotype of the type species of *Cylindroleberis* (type genus of the family), *Cypridina mariae* Baird, 1850 does not exist. In order to facilitate a systematic revision, this report lists the taxa and comments on taxonomic uncertainty, such as missing holotypes, description being only from a male (important diagnostic features are based on the female), or a description being from a unique specimen.

To begin, we review the taxonomic history of the Cylindroleberididae, followed by a summary of the higher systematics of the family as now accepted by Kornicker (e.g., 1981b); there are no current alternate views on this classification. Genera and their species are then listed alphabetically with necessary comments. Finally, a combined alphabetical list is provided of genera, species, and synonyms.

Taxonomic history of Cylindroleberididae

Cylindroleberidid ostracods were first described under the generic names *Cypridina* Milne-Edwards, 1840 and *Asterope* Philippi, 1840. Brady (1868) erected the genus *Cylindroleberis* but later (1871) synonymized it with *Asterope*. The subsequent

use of these genus names for the same generic concept caused some confusion. Müller used alternately *Asterope* (1890, 1912) and *Cylindroleberis* (1893, 1894, 1906). *Cylindroleberis* was preferred by some workers to avoid confusion with an echinoderm genus also named *Asterope* in 1840. However, some later authors (e.g., Skogsberg, 1920) continued to use *Asterope*, or later, another replacement name, *Asteropina* Strand, 1928.

By the late 19th century, cylindroleberidids were recognized as a group separate from cypridinids (Brady, 1871; Claus, 1876; Sars, 1865). Brady and Norman (1896) erected the family Asteropidae for *Asterope*. Brady (1897) erected a new genus *Cyclasterope*, which Müller (1906) referred with *Cylindroleberis* (=*Asterope*) to the Cylindroleberidinae, a new subfamily erected by him within the Cypridinidae.

The first major taxonomic work on the Cylindroleberididae was by Skogsberg (1920), part of a larger work on marine Ostracoda. Noting that the work to date on ostracods was largely based on missing material (e.g., Sars, 1887), or described the carapace only (e.g., Brady 1902) or considered too few characters, Skogsberg provided a thorough taxonomic treatment of the family. *Asterope* (=*Cylindroleberis*) and *Cyclasterope* were rediagnosed; *Cycloleberis* and *Asteropteron* were erected. Six new species were described and some existing species were given supplementary descriptions. Important diagnostic characters were established, such as the configuration of the s-seta, presence of the d-seta (on the first antenna) and e-seta (on the mandible).

Poulsen's (1965) *Dana Report* advanced ostracod taxonomy. The four existing cylindroleberidid genera were retained, and ten new ones erected, although six were *nomina nuda*. Poulsen did not designate type species for the new genera and thus only the monotypic genera (*Microasteropteron*, *Dolasterope*, *Bruuniella*, and *Archasterope*) were valid according to the International Code of Zoological Nomenclature. Five of Poulsen's remaining genera (*Asteropella*, *Heptonema*, *Diasterope*, *Parasterope*, and *Synasterope*) were later validated by Kornicker (1975) as discussed in Kornicker (1986). The genus *Philippiella* was not validated and its two species were later placed into other genera. To accommodate the expanding number of cylindroleberidid genera, Poulsen designated two subfamilies: the Cyclasteropinae (*Cyclasterope*, *Cycloleberis*,

Asteropteron and two new genera) and the Asteropinae (*Asteropina* (=*Cylindroleberis*) and eight new genera). These subfamilies were distinguished by setal counts on part of the first antenna and the maxilla endopod.

Most of the work on the Cylindroleberididae in the last fifty years is by Kornicker (e.g., 1974b, 1975, 1981b, 1986), including the description of more than one hundred new species and 24 genera. Kornicker (1981b) discussed the taxonomic history of the family.

Hartmann (1974) did not follow Poulsen's classification, instead erecting two new subfamilies: Cylindroleberidinae and Cycloleberidinae. Except for the placement of *Cyclasterope*, these subfamilies were the same as Poulsen's Asteropinae and Cyclasteropinae respectively. Kornicker (1981b) rejected this classification and retained that of Poulsen. He also erected the subfamily Asteropteroninae (1981b), split from Cyclasteropinae on the basis of carapace morphology. He adopted the subfamily name Cylindroleberidinae to reflect the effective replacement of the type genus *Asterope* by *Cylindroleberis*.

The type species of *Cylindroleberis* is *Cypridina mariae* Baird, 1850, subsequently designated by Sylvester-Bradley (1961). Unfortunately, the appendages were not described and the material is lost, leading to some uncertainty about the systematics of the group (Bowman and Kornicker, 1967).

Higher systematics of Cylindroleberididae

Family Cylindroleberididae Müller, 1906

Asteropidae Brady and Norman, 1896: 628

Cylindroleberidinae Müller, 1906: 32

Asteropinae.—Müller, 1912: 42

Cylindroleberididae.—Sylvester-Bradley, 1961: Q402 [new status]

Remarks. Although Brady and Norman created the family in 1896, it was based on a preoccupied name (*Asterope*). Hence the family is based on the earliest available name (*Cylindroleberis*). Poulsen (1965) split the family into two subfamilies, Asteropinae (=current Cylindroleberidinae) and Cyclasteropinae. The latter was further split by Kornicker (1981b) to make a new subfamily Asteropteroninae.

Subfamily Cylindroleberidinae Müller, 1906

Cylindroleberidinae Müller, 1906: 32

Asteropinae.—Poulsen 1965: 167

Remarks. Two tribes were introduced by Kornicker and Harrison-Nelson (2005) to separate the distinctive genus *Bruuniella* from the other genera.

Tribe Cylindroleberidini Müller, 1906

Cylindroleberidini.—Kornicker and Harrison-Nelson, 2005: 324 [new status]

Included genera. *Synasterope*, *Polyleberis*, *Cylindroleberis*, *Prionotoleberis*, *Postasterope*, *Homasterope*, *Monasterope*, *Parasterope*, *Heptonema*, *Diasterope*, *Xenoleberis*, *Bathyleberis*, *Domromeus*, *Skogsbergiella*, *Xandarasterope*, *Dolasterope*, *Archasterope*.

Tribe Bruuniellini Kornicker and Harrison-Nelson 2005

Bruuniellini Kornicker and Harrison-Nelson, 2005: 330.

Included genus: *Bruuniella*.

Subfamily Cyclasteropinae Poulsen, 1965

Cyclasteropinae Poulsen, 1965: 167.

Remarks. Kornicker (1981b) recognized three tribes. Hartmann's (1974) subfamily Cycloleberidinae was given new status as a tribe (although only retaining one of his genera, *Cycloleberis*, and two new genera). Cyclasteropini contained *Cyclasterope*, and a new tribe was erected, the Tetraleberidini, for two new genera.

Tribe Cyclasteropini Poulsen, 1965

Cyclasteropini.—Kornicker, 1981b: 73 [new status]

Included genus: *Cyclasterope*.

Tribe Tetraleberidini Kornicker, 1981

Tetraleberidini Kornicker, 1981b: 133.

Included genera: *Tetraleberis*, *Amboleberis*.

Tribe Cycloleberidini Hartmann, 1974

Cycloleberidinae Hartmann, 1974: [part; *Cycloleberis* only].

Cycloleberidini.—Kornicker, 1981b: 78 [new status].

Included genera: *Cycloleberis*, *Leuroleberis*, *Alphaleberis*.

Subfamily Asteropteroninae Kornicker, 1981

Asteropteroninae Kornicker, 1981b: 189.

Included genera: *Asteropella*, *Omegasterope*, *Pteromeniscus*, *Macroasteropteron*, *Microasteropteron*, *Asteropteron*, *Actinoseta*, *Asteropterygion*.

Checklist (alphabetical by genus and species)

Actinoseta Kornicker, 1958

Actinoseta Kornicker, 1958: 243

Type species. *Actinoseta chelisparsa* Kornicker, 1958: 244 by original designation.

Actinoseta chelisparsa Kornicker, 1958

Actinoseta chelisparsa Kornicker, 1958: 244

Type locality. Bimini, Bahamas, USA.

Actinoseta hummelincki Kornicker, 1981

Actinoseta hummelincki Kornicker, 1981b: 201

Type locality. Virginia Key, Florida, USA, 2 m depth.

Actinoseta jonesi Kornicker, 1981

Actinoseta jonesi Kornicker, 1981b: 210

Type locality. West of Punta Charagata, Isla Cubaque, Venezuela, <1.5 m depth.

Actinoseta nodosa Kornicker, 1981

Actinoseta nodosa Kornicker, 1981b: 215

Type locality. Male Atoll, Maldives, Indian Ocean, 0.3 m depth.

Alphaleberis Kornicker, 1981

Alphaleberis Kornicker, 1981b: 123

Type species. *Alphaleberis alphathrix* Kornicker, 1981 by original designation.

Alphaleberis alphathrix Kornicker, 1981

Alphaleberis alphathrix Kornicker, 1981b: 123

Type locality. East Beach, Nosy Ve Island, Madagascar, infralittoral.

Amboleberis* Kornicker, 1981Amboleberis* Kornicker, 1981b: 167

Type species. *Asterope americana* Müller, 1890 by original designation.

***Amboleberis americana* (Müller, 1890)**

- Asterope americana* Müller, 1890: 240
Asterope dentata.—Müller, 1890: 240 [lapsus]
Cylindroleberis americana.—Müller, 1894: 218 [inferred]
Cyclasterope brevis.—Brady, 1902: 183
Cyclasterope americana.—Müller, 1912: 33
Cyclasterope americana.—Müller, 1912: 48
Cyclasterope deutata.—Müller, 1912: 48 [misspelling of *C. dentata* in synonymy]
Cycloleberis americana.—Skogsberg, 1920: 442
Cyclasterope priacanthus Tressler, 1949: 338
Cyclasterope tripla Tressler, 1949: 340
Cyclasterope sphaerica Tressler, 1949: 336
Cycloleberis biminiensis Kornicker, 1958: 232, 243
Cyclasterope biminiensis.—Darby, 1965: 30
Cycloleberis priacanthus.—Poulsen, 1965: 245
Cycloleberis tripla.—Poulsen, 1965: 245
Cycloleberis sphaerica.—Poulsen, 1965: 245
Amboleberis americana.—Kornicker, 1981b: 168

Type locality. Coast of South America (Pernambuco), pelagic.

Taxonomic decision for synonymy. Kornicker (1981b: 168) [most recent synonym list]

Remarks. The holotype is likely to be lost. A supplementary description of an adult female was provided by Kornicker (1981b).

Amboleberis antyx* Kornicker, 1981Amboleberis antyx* Kornicker, 1981b: 181

Type locality. outer slope, Grand Recif, Madagascar, 17 m depth.

Amboleberis cubensis* Lalana and Kornicker, 1997Amboleberis cubensis* Lalana and Kornicker, 1997: 108

Type locality. Cayo los Dromedarios, north coast of Cuba, 1 m depth.

Remarks. The species is based on a unique specimen.

Archasterope* Poulsen, 1965Archasterope* Poulsen, 1965: 339

Philippiella Poulsen, 1965: 344 [part] [nomen nudum, without type species; 2 included species, *Asterope quinquesetae* now included in *Archasterope*, and *Asterope spinifera* now in *Skogsbergiella*]

Empoulsenia Kornicker, 1975: 499

Type species. *Archasterope dentata* Poulsen, 1965 by monotypy.

Taxonomic decision for synonymy. Some species of *Empoulsenia* (including the type species *Empoulsenia quinquesetae*) were included in *Archasterope* by Kornicker (1993); synonymy by Kornicker (in Kornicker and Poore, 1996: 142).

Remarks. The genus diagnosis was emended by Kornicker (in Kornicker and Poore, 1996: 142).

Archasterope altrix* Kornicker, 1996Archasterope altrix* Kornicker, in Kornicker and Poore, 1996: 154

Type locality. South of Point Hicks, Victoria, Australia, 800 m depth. 38°16.40'S, 149°27.60'E.

***Archasterope antarctica* (Kornicker, 1975)**

Empoulsenia antarctica Kornicker, 1975: 509
Archasterope antarctica.—Kornicker, 1993: 116

Type locality. Weddell Sea, South Shetland Islands, Antarctic, 220–240 m depth.

Archasterope apex* Kornicker, 1996Archasterope apex* Kornicker, in Kornicker and Poore, 1996: 150

Type locality. off Freycinet Peninsula, Tasmania, Australia, 500 m depth. 47°58.60'S, 148°38.80'E.

Remarks. The species is based on a unique specimen.

Archasterope bulla* Kornicker, 1975Archasterope bulla* Kornicker, 1975: 384

Type locality. Tierra del Fuego, Strait of Magellan, Subantarctic, 280 m depth.

Archasterope dentata* Poulsen, 1965Archasterope dentata* Poulsen, 1965: 339

Type locality. off southeast Australia, 180 m depth. 38°12'S, 149°40'E.

Remarks. The first antennae are missing from the holotype and paratype (Kornicker, 1989). Thus, it cannot be verified whether the d-seta is a filament or a spine. Other species in the genus have a filament.

Archasterope efficax* Kornicker, 1996Archasterope efficax* Kornicker, in Kornicker and Poore, 1996: 143

Type locality. off Freycinet Peninsula, Tasmania, Australia, 400 m depth. 41°57.50'S, 148°37.90'E.

***Archasterope monothrix* (Kornicker, 1988)**

Empoulsenia monothrix Kornicker, 1988: 24
Archasterope monothrix.—Kornicker, in Kornicker and Poore, 1996: 142

Type locality. Beaufort Sea, Arctic Ocean, 101 m depth. 71°23'30" N, 152°43'00" W.

***Archasterope pentathrix* (Kornicker, 1971)**

Philippiella pentathrix Kornicker, 1971: 205
Empoulsenia pentathrix.—Kornicker, 1975: 502
Archasterope pentathrix.—Kornicker, 1993: 116

Type locality. Greenwich Island, South Shetland Islands, Antarctica.

Archasterope polythrix* (Chavtur, 1983)Archasterope polythrix* Chavtur, 1983 [reference not sighted]

Remarks. We were unable to obtain this reference; hence we do not have information about the type locality.

***Archasterope quinquesetae* (Skogsberg, 1920)**

Asterope quinquesetae Skogsberg, 1920: 470
Philippiella quinquesetae.—Poulsen, 1965: 344
Empoulsenia quinquesetae.—Kornicker, 1975: 501
Archasterope quinquesetae.—Kornicker, 1993: 112

Type locality. South Georgia, off mouth of Cumberland Bay. 54°11'S, 36°18'W.

***Archasterope unisetosa* (Hartmann, 1989)**

Empoulsenia unisetosa Hartmann, 1989: 236
Archasterope unisetosa.—Kornicker, 1993: 119

Type locality. vicinity of Elephant Island, South Georgia district, Antarctic region, 427 m depth.

***Archasterope verax* Kornicker, 1996**

Archasterope verax Kornicker, in Kornicker and Poore, 1996: 157

Type locality. off Freycinet Peninsula, Tasmania, Australia, 1264 m depth. 42°00.25'S, 148°43.55'E.

***Archasterope weddellensis* (Kornicker, 1975)**

Empoulsenia weddellensis Kornicker, 1975: 514
Archasterope weddellensis.—Kornicker, 1993: 117

Type locality. Continental subregion, Antarctic region, 132-650 m depth.

Remarks. A juvenile (A-1) female is known, but no adult female is known.

***Asteropella* Kornicker, 1975**

Asteropella Poulsen, 1965: 222 [nomen nudum; without type species; 3 included species, *Asteropella mortenseni*, *A. punctata*, *A. monambon*]
Asteropella Kornicker, 1975: 557 [designated type species]

Type species. *Asteropella mortenseni* Poulsen, 1965 by original designation, Kornicker, 1975.

***Asteropella agassizii* (Müller, 1870)**

Cypridina Agassizii Müller, 1870: 255
Cylindroleberis Agassizii.—Müller, 1894: 218. [combination inferred]
Cyclasterope agassizii.—Müller, 1912: 48
Asteropteron Agassizi.—Skogsberg, 1920: 88
Cyclasterope agassizi.—Poulsen, 1965: 225
Asteropella agassizii.—Kornicker, 1981b: 231

Type locality. Florianopolis, Santa Catarina Island, southern coast of Brazil.

Remarks. There is no holotype selected.

***Asteropella kalkei* Kornicker, 1986**

Asteropella kalkei Kornicker, 1986: 117

Type locality. Off Weeks Island, Louisiana, USA, 6.4 m depth. 29°04'22"N, 91°44'24"W.

***Asteropella kaufmani* Kornicker, 1981**

Asteropella kaufmani Kornicker, 1981b: 268

Type locality. Naos Island, Panama, 1.2 m depth.

***Asteropella maclaughlinae* Kornicker, 1981**

Asteropella maclaughlinae Kornicker, 1981b: 244

Type locality. Anclote Anchorage, west coast of Florida, USA, 1 m depth.

***Asteropella monambon* (Kornicker, 1958)**

Asteropteron monambon Kornicker, 1958: 281
Asteropella monamba.—Poulsen 1965: 225
Asteropella monambon.—Kornicker, 1975: 561

Type locality. Bimini, Bahamas, 3-20 m depth.

Remarks. The sex of the holotype was unknown at the time of description. A subsequent analysis by Kornicker (1981b) indicated that the holotype was a male. A supplementary description of the adult male and a description of the adult female were provided by Kornicker (1981b).

***Asteropella mortenseni* Poulsen, 1965**

Asteropella mortenseni Poulsen, 1965: 169

Type locality. Virgin Islands, West Indies, 20-40 m depth.

***Asteropella pax* Kornicker, 1986**

Asteropella species 1 Kornicker, 1981b: 275
Asteropella pax Kornicker, 1986: 124

Type locality. Gulf of Mexico, 15 m depth.

***Asteropella punctata* Poulsen, 1965**

Asteropella punctata Poulsen, 1965: 169
Asteropella punctatum Poulsen, 1965: 478

Type locality. East coast of Thatch Island, West Indies, 25-30 m depth.

***Asteropella rotundicostata* (Hartmann, 1965)**

Cycloleberis rotundicostata Hartmann, in Hartmann-Schröder and Hartmann, 1965: 326
Asteropella rotundicostata.—Kornicker, 1975: 559

Type locality. Bahia Ingles off Chile, 41°48'S, 75°53'W.

Remarks. The species is based on a unique specimen.

***Asteropella scammonensis* McKenzie, 1965**

Asteropella scammonensis McKenzie, 1965: 60

Type locality. Scammon lagoon, Baja California.

Remarks. Kornicker (1981b: 40) considers the species description to be based on juvenile specimens.

***Asteropella slatteryi* Kornicker, 1981**

Asteropella species Tuel et al., 1976: 141
Asteropella slatteryi Kornicker, 1981b: 260

Type locality. Northern part of Monterey Bay, California, USA, 16 m depth. 36°57'06"N, 121°56'12"W.

***Asteropella trithrix* Kornicker, 1981**

Asteropella trithrix Kornicker, 1981b: 251

Type locality. Bahia de los Angeles, 7 m depth.

***Asteropteron* Skogsberg, 1920**

Asteropteron Skogsberg, 1920: 442

Type species. *Asterope fusca* Müller, 1890 by original designation.

***Asteropteron fuscum* (Müller, 1890)**

Asterope fusca Müller, 1890: 242
Cylindroleberis fusca.—Kajiyama, 1912: 617
Cyclasterope fusca.—Müller, 1912: 48
Asteropteron fuscum.—Skogsberg, 1920: 443

Type locality. off the Japanese coast, Enoshima, Fujisawa-shi.

Remarks. There are five syntypes for this species. Hiruta (1979) redescribed the species based on new specimens from Kyushu, Japan. Kornicker (1981b) discussed and redescribed Müller's type specimens.

***Astropterygion* Kornicker, 1981**

Astropterygion Kornicker, 1981b: 281

Type species. *Astropterygion thomassini* Kornicker, 1981 by original designation.

***Astropterygion climax* Kornicker, 1996**

Astropterygion climax Kornicker, 1996: 65

Type locality. Gulf of Carpentaria, in vicinity of Weipa, Queensland, Australia, 10 m depth.

Remarks. The species is based on male specimens only; there are no known females.

***Asteropterygion dayi* Kornicker, 1981**

Asteropterygion dayi Kornicker, 1981b: 321

Type locality. Langebaan Lagoon, South Africa, 4.5 m depth.

***Asteropterygion hirsutum* (Poulsen, 1965)**

Asteropteron hirsutum Poulsen, 1965: 187

Asteropterygion hirsutum.—Kornicker, 1981: 321

Type locality. Malayan Archipelago, Danish Key Island Expedition, 30 m depth.

Remarks. The species is based on a unique specimen which is a juvenile male.

***Asteropterygion hulingsi* (Kornicker, 1975)**

Asteropteron hulingsi Kornicker, 1975: 562

Asteropterygion hulingsi.—Kornicker, 1981b: 289

Type locality. South Atlantic, east of Argentina, 44 m depth. 40°11'S, 60°27'W.

Remarks. There are no adult specimens known for this species.

***Asteropterygion liguriae* (Granata, 1915)**

Cyclasterope liguriae Granata, 1915: 30

Asteropterygion liguriae.—Kornicker, 1981b: 284

Type locality. Bay of Thursday Island, Torres Strait, 4 m depth. 10°35'08"S, 142°13'14"E.

Remarks. The specimens could be conspecific with *Asteropterygion magnum* (Kornicker, 1981b). The holotype is probably lost.

***Asteropterygion magnum* (Poulsen, 1965)**

Asteropteron magnum Poulsen, 1965: 178

Asteropterygion magnum.—Kornicker, 1981b: 320

Type locality. Off Disaster Bay, southeast Australia, 65-90 m depth.

***Asteropterygion nodulosum* (Poulsen, 1965)**

Asteropteron nodulosum Poulsen, 1965: 201

Asteropterygion nodulosum.—Kornicker, 1981b: 286

Type locality. False Bay, Cape of Good Hope, South Africa, 20 m depth.

Remarks. The species is based on an adult male only, there are no known adult females for this species.

***Asteropterygion oculitristis* (Darby, 1965)**

Asteropteron oculitristis Darby, 1965: 29

Asteropterygion oculitristis.—Kornicker, 1981b: 290

Type locality. Off Sapelo Island, Georgia, USA, 18 m depth.

***Asteropterygion peterseni* Kornicker, 1981**

Asteropterygion peterseni Kornicker, 1981b: 328

Type locality. Off Tanzania, in vicinity of Kunduchi, 0-25 m depth.

***Asteropterygion romei* Kornicker, 1981**

Asteropterygion romei Kornicker, 1981b: 312

Type locality. Off Kunduchi, Tanzania, 2 m depth.

***Asteropterygion samoae* Kornicker and Harrison-Nelson, 2006**

Asteropterygion samoae Kornicker and Harrison-Nelson, 2006: 252

Type locality. Fagatele Bay, Tutuila, American Samoa, 21-24 m depth. 14°21.96'S, 170°45.85'W.

Remarks. The species is based on a unique juvenile female specimen.

***Asteropterygion setiferum* (Kornicker and Caraion, 1974)**

Asteropteron setiferum Kornicker and Caraion, 1974: 66

Asteropterygion setiferum.—Kornicker, 1981b: 286

Type locality. Off Mauritania, 1100? m depth.

***Asteropterygion skogsbergii* (Poulsen, 1965)**

Asteropteron skogsbergii Poulsen, 1965: 206

Asteropteron cf. spinosum Hartmann, 1974: 235

Asteropterygion skogsbergii.—Kornicker, 1981b: 299

Type locality. off Mombasa, Kenya, 33 m depth.

Taxonomic decision for synonymy. Kornicker (1981b).

Remarks. The labelling of the holotype is uncertain, (Kornicker, 1981b: 300).

***Asteropterygion spinosum* (Poulsen, 1965)**

Asteropteron spinosum Poulsen 1965: 212

Asteropterygion spinosum.—Kornicker, 1981b: 300

Type locality. Off Mombasa, Kenya, 33 m depth.

Remarks. The holotype is considered lost. The species is based on a unique specimen (Kornicker, 1981b).

***Asteropterygion thailandicum* (Poulsen, 1965)**

Asteropteron thailandicum Poulsen, 1965: 193

Asteropterygion thailandicum.—Kornicker, 1981b: 286

Type locality. Koh Kam, Thailand, 18 m depth.

Remarks. There are no adult females known for this species.

***Asteropterygion thomassini* Kornicker, 1981**

Asteropterygion thomassini Kornicker, 1981b: 301

Type locality. Tulear Lagoon, Southern Pass, Madagascar.

***Bathyleberis* Kornicker, 1975**

Bathyleberis Kornicker, 1975: 538

Type species. *Bathyleberis grossmani* Kornicker, 1975 by original designation.

***Bathyleberis babax* Kornicker, 1996**

Bathyleberis babax Kornicker, in Kornicker and Poore, 1996: 173

Type locality. South of Point Hicks, Victoria, Australia, 1500 m depth. 38°25.00'S, 149°00.00'E.

***Bathyleberis garthi* Baker, 1979**

Bathyleberis garthi Baker, 1979: 293

Type locality. off Sand Point, California, USA, 96 m depth. 34°13'17"N, 119°29'15"W.

***Bathyleberis grossmani* Kornicker, 1975**

Bathyleberis grossmani Kornicker, 1975: 539

Type locality. Drake Passage, Antarctic, 3678-3816 m depth.

Bathyleberis hancocki* Baker, 1979Bathyleberis hancocki* Baker, 1979: 297

Type locality. Point Dune, California, USA, 33°59'10" N, 118°48'15" W.

Bathyleberis kurilensis* (Chavtur, 1978)Empoulsenia kurilensis* Chavtur, 1978: 154*Bathyleberis kurilensis*.—Kornicker, 1986: 62

Type locality. Sea of Okhotsk coast of Urop Island, 20 m depth.

Remarks. An emended diagnosis was provided by Chavtur (1983: 81).

Bathyleberis monothrix* Kornicker, 1975Bathyleberis monothrix* Kornicker, 1975: 545

Type locality. South Atlantic, East of Argentina, 82 m depth. 41°27'S, 59°33'W.

Bathyleberis oculata* Kornicker, 1975Bathyleberis oculata* Kornicker, 1975: 551

Type locality. Cook Strait, New Zealand, 117 m depth. 40°33'S, 174°07'E.

Bathyleberis thrix* Kornicker, 1988Bathyleberis thrix* Kornicker, 1988: 28

Type locality. Beaufort Sea, Arctic Ocean, 53 m depth. 71°17'54"N, 154°33'130'W.

Bathyleberis toxotes* Kornicker, 1986Bathyleberis toxotes* Kornicker, 1986: 63

Type locality. Gulf of Mexico, off Galveston, Texas, 54 m depth.

Bruuniella Poulsen, 1965*Bruuniella Poulsen, 1965: 323* [no diagnosis provided]

Type species. *Bruuniella breviata* Poulsen, 1965: 323 by monotypy.

Remarks. A diagnosis was not provided by Poulsen (1965). Kornicker and Harrison-Nelson (2005: 330) created a new tribe for the genus (*Bruuniellini*) and provided a diagnosis.

Bruuniella alpha* Kornicker and Harrison-Nelson, 2005Bruuniella species A* Kornicker, 1981a: 12*Bruuniella alpha* Kornicker and Harrison-Nelson, 2005: 334

Type locality. Smith's Sound, Governor's Island, Bermuda, 0.6 m depth.

Remarks. The species is known only from one specimen, a juvenile male, which could be conspecific with *Bruuniella hazeli* (Kornicker and Harrison-Nelson, 2005: 335).

Bruuniella beta* Kornicker and Harrison-Nelson, 2005Bruuniella beta* Kornicker and Harrison-Nelson, 2005: 335

Type locality. Johnston Atoll, 16°44'52.3"N, 169°31'3.5"W, 2-9.5 m depth.

Bruuniella breviata* Poulsen, 1965Bruuniella breviata* Poulsen, 1965: 323

Type locality. Koh Chang, Thailand, 2 m depth.

Remarks. A supplementary description was provided by Kornicker and Harrison-Nelson (2005: 333).

Bruuniella hazeli* Kornicker, 1986Bruuniella hazeli* Kornicker, 1986: 85

Type locality. Vineyard Haven Harbor, Martha's Vineyard, Massachusetts, 1.5 m depth.

Remarks. A supplementary description and a discussion of the adult female holotype was provided by Kornicker and Harrison-Nelson (2005: 333).

Cyclasterope* Brady, 1897Cyclasterope* Brady, 1897

Type species. *Cyclasterope hendersoni* Brady, 1897, by subsequent designation (Skogsberg, 1920: 441).

Remarks. An emended diagnosis was provided by Skogsberg (1920). A supplementary diagnosis was provided by Poulsen (1965: 283).

Cyclasterope arthuri* (Stebbing, 1900)Asterope* species Stebbing, 1899: 31*Asterope arthuri* Stebbing, 1900: 607*Cylindroleberis hilgendorfii* Müller, 1906: 33 [part]*Asterope hilgendorfii*.—Müller, 1912: 44 [part]*Cyclasterope arthuri*.—Kornicker, 1981b: 77

Type locality. Blanche Bay, New Britain, Bismark Archipelago, 110-128 m.

Taxonomic decision for synonymy. Kornicker (1981b) [most recent list].

Remarks. There was no holotype selected for this species. Because no female appendages were drawn or described, "additional information concerning this species is needed to be certain that it is valid" (Kornicker, 1981b: 77).

Cyclasterope bisetosa* Poulsen, 1965Cyclasterope bisetosa* Poulsen, 1965: 301

Type locality. Padang Harbour, west coast of Sumatra, <50 m depth.

Remarks. This species is based on one female and one juvenile male only.

Cyclasterope fascigera* Brady, 1902Cyclasterope fascigera* Brady, 1902: 181*Cylindroleberis fascigera*.—Müller, 1906: 33*Asterope fascigera*.—Müller, 1912: 43*Cycloleberis fascigera*.—Poulsen, 1965: 476 [obvious lapsus]

Type locality. Java.

Taxonomic decision for synonymy. Kornicker (1981b) [most recent list].

Remarks. There was no holotype selected. The species was redescribed by Kornicker (1981b).

Cyclasterope hendersoni* Brady, 1897Cyclasterope hendersoni* Brady, 1897: 86*Cylindroleberis hilgendorfii* Müller, 1906: 33 [part]*Asterope hilgendorfii* Müller, 1912: 44 [part]

Type locality. Madras Harbour, India, 7.3-9.1 m depth.

Taxonomic decision for synonymy. Kornicker, 1981b [most recent list].

Remarks. There was no holotype selected.

***Cyclasterope hilgendorfii* (Müller, 1890)**

Asterope hilgendorfii Müller, 1890: 221
Cylindroleberis hilgendorfii.—Müller, 1894: 63
Cyclasterope hilgendorfii.—Hanai, 1959: 425

Type locality. Coast of Japan, Enosima (Shonan), 22 m depth.

Remarks. The species is based on a unique specimen. See Kornicker (1981b) for remarks about the identity of specimens.

***Cyclasterope albomaculata* (Baird, 1860)**

Cypridina albo-maculata Baird, 1860: 201
Cypridina albomaculata.—Brady, 1897: 87
?Cyclasterope albomaculata.—Kornicker, 1975: 30

Type locality. Swan River, near Perth, Western Australia.

Remarks. The type specimen (or new specimens) need examination for clarification of the generic placement, and it is possible that these specimens are conspecific with *Leuroleberis mackenziei* (Kornicker, 1981b: 78).

Cycloleberis Skogsberg, 1920

Cycloleberis Skogsberg, 1920

Type species. *Cylindroleberis lobiancoi* Müller, 1894 by original designation.

***Cycloleberis christiei* Kornicker and Maddocks, 1977**

Cycloleberis galatheae Poulsen, 1965: 260 [part; juvenile male only]
Cycloleberis christiei Kornicker and Maddocks, 1977: 894

Type locality. Langebaan Lagoon, South Africa. 33°07'S, 18°01'E

Taxonomic decision for synonymy. Kornicker and Maddocks (1977).

***Cycloleberis galatheae* Poulsen, 1965**

Cycloleberis galatheae Poulsen, 1965: 260 [part; juvenile A-1 female holotype].

Type locality. False Bay, South Africa, 20 m depth.

Remarks. The holotype is based on a juvenile (A-1) female. Kornicker (1981b: 80) believes that *Cycloleberis galatheae* may be conspecific with *Cycloleberis squamiger*.

***Cycloleberis lobiancoi* (Müller, 1893)**

Cylindroleberis Lobianci Müller, 1893: 359
Cyclasterope lobianci.—Müller, 1906: 33
Cyclasterope lobiancoi.—Müller, 1912: 48 [part]
Cycloleberis Lobianci.—Skogsberg, 1920: 437
Cycloleberis lobiancoi.—Poulsen, 1965: 238
Cyclasterope (Cycloleberis) lobiancoi.—Sylvester-Bradley, 1961: Q402

Type locality. Gulf of Naples.

Taxonomic decision for synonymy. Kornicker (1981b) [most recent list].

Remarks. There was no holotype selected. The original specimens are considered lost (Kornicker, 1974b). Description of another specimen was provided by Kornicker (1974b).

***Cycloleberis squamiger* (Scott, 1894)**

Asterope squamiger Scott, 1894: 140
Cycloleberis squamiger.—Kornicker and Caraion, 1974: 47

Type locality. Sao Tome Island, Gulf of Guinea.

Remarks. A lectotype is available by subsequent designation (Kornicker and Caraion, 1974).

***Cylindroleberis* Brady, 1868**

Asterope Philippi, 1840: 186
Cylindroleberis Brady, 1868: 127 [replacement name; *Asterope* preoccupied]
Asteropina Strand, 1928 [unnecessary replacement name; *Asterope* preoccupied]

Type species. *Cylindroleberis mariae* (Baird, 1850) by subsequent designation Sylvester-Bradley, 1961.

Remarks. The genus was redescribed by Skogsberg (1920: 445) (using *Asterope*) and Poulsen (1965: 434) (using *Asteropina*). The composition of the genus was discussed by Kornicker (1991a: 110).

***Cylindroleberis bacescui* Kornicker and Caraion, 1974**

Cylindroleberis bacescui Kornicker and Caraion, 1974: 29

Type locality. Mauritania, 30 m depth. 20°00'07"N, 17°18'04"W.

***Cylindroleberis grimaldi* (Skogsberg, 1920)**

Asterope grimaldi Skogsberg, 1920: 510
Asteropina grimaldi.—Poulsen, 1965: 434
Cylindroleberis grimaldi.—Kornicker, 1991a: 110

Type locality. Mediterranean, Monaco harbour, 10 m depth.

***Cylindroleberis grimaldi vicina* (Skogsberg, 1920)**

Cylindroleberis oblonga Müller 1894: 219
Asterope mariae Müller 1912: 45 [part]
Asterope grimaldi var. *vicina* Skogsberg, 1920: 518
Asteropina grimaldi var. *vicina*.—Poulsen, 1965: 434
Cylindroleberis grimaldi var. *vicina*.—Kornicker, 1991a: 110

Type locality. Villefranche sur mer, France, Mediterranean sea, 95 m depth.

Taxonomic decision for synonymy. Skogsberg (1920).

***Cylindroleberis kliei* Kornicker, 1976**

Cylindroleberis kliei Kornicker, 1976b: 2

Type locality. Luderiz Bay, South-west Africa.

***Cylindroleberis mariae* (Baird, 1850)**

Cypridina mariae Baird, 1850: 257
Cylindroleberis mariae.—Brady, 1868
Asterope mariae.—Brady, 1871
Asteropina mariae.—Strand, 1928

Type locality. Off Isle of Skye, Scotland, UK.

Remarks. Baird described only the carapace of this species. A holotype was not designated nor deposited, and is considered lost (Skogsberg, 1920). See Skogsberg (1920: 518) for a discussion of specimens.

***Cylindroleberis marranyin* Syme and Poore, 2006**

Cylindroleberis marranyin Syme and Poore, 2006: 61

Type locality. Sandy Point, Western Port, Victoria, Australia. 38° 24' S, 145° 14' E, benthic, 4 m.

***Cylindroleberis minuta* (Poulsen, 1965)**

Asteropina minuta Poulsen, 1965: 436
Cylindroleberis minuta.—Kornicker, 1991a: 110

Type locality. Koh Kam, Thailand, 9 m depth.

***Cylindroleberis nodulifera* (Poulsen, 1965)**

Asteropina nodulifera Poulsen, 1965: 447
Cylindroleberis nodulifera.—Kornicker, 1991a: 110

Type locality. Guardaga, Red Sea, surface at night.

Remarks. The species is known only from adult males.

***Cylindroleberis rangiroaensis* Hartmann, 1984**

Cylindroleberis rangiroaensis Hartmann, 1984: 119

Type locality. Kalkalgen vom Anleger der Kia Ora Village Lagune.

Remarks. This species has a long mandibular exopod and thus it does not fit the genus *Cylindroleberis*; it should be moved (Kornicker, 1991a).

***Cylindroleberis thailandica* (Poulsen, 1965)**

Asteropina thaliandica Poulsen, 1965: 444

Cylindroleberis thailandica.—Kornicker, 1991a: 110

Type locality. Koh Kahdat, Thailand, 9 m depth.

Remarks. The species is known only from one adult female, one juvenile female, and one juvenile male.

***Cylindroleberis variabilis* Kornicker, 1970**

Cylindroleberis variabilis Kornicker, 1970: 19

Type locality. Sacol Island, Zamboanga Province, Phillipines.

***Cylindroleberis verrucosa* (Poulsen, 1965)**

Asteropina verrucosa Poulsen, 1965: 440

Cylindroleberis verrucosa.—Kornicker, 1991a: 110

Type locality. Koh Chang, Thailand, 2 m depth.

Remarks. The species is known only from one adult female and one juvenile male.

***Cylindroleberis vibex* Kornicker, 1992**

Cylindroleberis vibex Kornicker, 1992: 185

Type locality. Mozambique Channel, Mayotte, 6 m depth.

Remarks. A supplementary description of the adult female and description of adult male was provided by Kornicker (1998).

***Cylindroleberis vix* Kornicker, 1992**

Cylindroleberis vix Kornicker, 1992: 179

Type locality. Mozambique Channel, Mayotte, 15-20 m depth.

Remarks. A supplementary description of the adult female and a description of the adult male was provided by Kornicker and Thomassin (1998).

***Diasterope* Kornicker, 1975**

Diasterope Poulsen, 1965: 345 [nomen nudum; without type species; 4 included species, *Diasterope canina*, *D. tenuiseta*, *D. pilosa*, *D. bisetosa*].

Diasterope Kornicker, 1975: 388 [designated type species]

Type species. *Diasterope pilosa* Poulsen, 1965 by original designation Kornicker, 1975.

***Diasterope bisetosa* Poulsen, 1965**

Diasterope bisetosa Poulsen, 1965: 355

Type locality. Red Sea, surface.

Remarks. The species is known only from adult males.

***Diasterope canina* Poulsen, 1965**

Diasterope canina Poulsen, 1965: 358

Type locality. St. Croix, off Buck Island, Virgin Islands, West Indies, 7 m depth.

Remarks. The species is known only from juvenile males.

***Diasterope grisea* (Brady, 1898)**

Asterope grisea Brady 1898: 432

Cylindroleberis? grisea Eagar, 1971: 61

Diasterope grisea.—Kornicker, 1975: 389

Type locality. Akaroa Harbor, New Zealand, 11 m depth.

Remarks. There was no holotype designated. A supplementary description of the female was provided by Kornicker (1975).

***Diasterope persyiensis* Chavtur, 1983**

Diasterope persyiensis Chavtur, 1983 [reference not sighted]

Remarks. We were unable to obtain this reference; hence we do not have information about the type locality.

***Diasterope pilosa* Poulsen, 1965**

Diasterope pilosa Poulsen, 1965: 347

Type locality. Pacific coast of Canada, Strait of Georgia, Lasqueti Island, 45 m depth.

***Diasterope procax* Kornicker and Iliffe, 2000**

Diasterope procax Kornicker and Iliffe, 2000: 61

Type locality. Exuma Sound, Bahamas, 88 m depth.

***Diasterope schmitti* Kornicker, 1975**

Diasterope schmitti Kornicker, 1975: 395

Type locality. Robertson Bay, west of Cape Adare, Ross Sea, Antarctic, 128 m depth.

***Diasterope tenuiseta* Poulsen, 1965**

Diasterope tenuiseta Poulsen, 1965: 351

Type locality. North of Loanga, Virgin Islands, West Indies, 30 m depth.

Remarks. The species is known only from juvenile males.

***Diasterope wirraka* Syme and Poore, 2006**

Diasterope wirraka Syme and Poore, 2006: 56

Type locality. Eastern Bass Strait, Victoria, Australia. 15.3 km ESE of eastern edge of Lake Tyers. 37°53.23' S, 148°15.24' E, benthic, 43 m.

***Dolasterope* Poulsen, 1965**

Dolasterope Poulsen, 1965: 317

Type species. *Dolasterope johanseni* Poulsen, 1965 by monotypy.

***Dolasterope johanseni* Poulsen, 1965**

Dolasterope johanseni Poulsen, 1965: 318

Type locality. entrance to Milford Sound, New Zealand, 136 m depth. 44°34'S 167°48'E.

Remarks. The species is based on a unique specimen.

***Domromeus* Kornicker, 1989**

Domromeus Kornicker, 1989: 124

Type species. *Domromeus heptathrix* Kornicker, 1989 by original designation.

Remarks. An emended genus diagnosis was provided by Kornicker (in Kornicker and Poore, 1996: 170).

Domromeus heptathrix* Kornicker, 1989Domromeus heptathrix* Kornicker, 1989: 124*Type locality.* west of Bay of Biscay, 2190 m depth.
47°33'24"S 8°40'42"W.***Dromromeus merx* Kornicker, 1996***Dromromeus merx* Kornicker, in Kornicker and Poore, 1996: 170*Type locality.* off Eden, New South Wales, Australia, 363 m depth. 37°00.60"S 150°20.70"E.*Remarks.* The species is based on a unique specimen, a juvenile (A-1) male. We believe this specimen to be very similar to *Bathyleberis grossmani*.***Heptonema* Cohen and Kornicker, 1975***Heptonema* Poulsen 1965: 328 [nomen nudum; without type species; 2 included species, *Heptonema serrata*, *H. keiensis*]*Heptonema* Cohen and Kornicker, 1975: 23 [designated type species]*Type species.* *Heptonema serratum* Poulsen, 1965, by original designation, Kornicker, 1975.*Remarks.* An emended diagnosis was provided by Kornicker (1992).***Heptonema homelix* Kornicker, 1991***Heptonema homelix* Kornicker, 1991a: 118*Type locality.* Enewetak lagoon, midwater trawl.***Heptonema keiensis* Poulsen, 1965***Heptonema keiensis* Poulsen, 1965: 334*Type locality.* East Indies, Kei Islands, 25 m depth.*Remarks.* The species is known only from juveniles.***Heptonema latex* Kornicker, 1992***Heptonema latex* Kornicker, 1992: 172*Type locality.* Mozambique Channel, Mayotte, 32 m depth.***Heptonema latum* Kornicker, 1986***Heptonema latum* Kornicker, 1986: 54*Type locality.* Continental shelf, New Jersey, 110-112 m depth.***Heptonema serratum* Poulsen, 1965***Heptonema serrata* Poulsen, 1965: 329*Heptonema serratum*.—Kornicker, 1986: 54*Type locality.* West coast of Thatch Island, Virgin Islands, West Indies, 25-30 m depth.***Homasterope* Kornicker, 1975***Homasterope* Kornicker, 1975: 524*Type species.* *Asterope curta* Skogsberg, 1920 by original designation.***Homasterope curta* (Skogsberg, 1920)***Asterope curta* Skogsberg 1920: 498*Parasterope curta*.—Poulsen, 1965: 363*Homasterope curta*.—Kornicker, 1975: 525*Type locality.* South Georgia, Antarctic region, 24-52 m depth.
54°22'S, 36°27'W.***Homasterope glacialis* (Müller, 1908)***Cylindroleberis ovalis* Müller, 1908: 93*Asterope glacialis* Müller, 1912: 47 [new name]*Homasterope glacialis*.—Kornicker, 1975: 526*Type locality.* Gauss Station, Antarctica, 385 m depth.*Remarks.* This species is not well enough known to distinguish it from *H. curta* or *H. maccaini* (Kornicker, 1993).***Homasterope maccaini* Kornicker, 1975***Homasterope maccaini* Kornicker, 1975: 528*Type locality.* Arthur Harbour, Anvers Island. Antarctic region, 6-98 m depth.***Homasterope micra* Kornicker, 1975***Homasterope micra* Kornicker, 1993: 534*Type locality.* off Golfo de Penas, west of Chile, Antarctic, 2657 m depth.*Remarks.* Although no adult males are known, Kornicker placed this species into this genus on the basis of the number of seta on the third article of the first antenna.***Homasterope trebax* Kornicker, 1996***Homasterope trebax* Kornicker, in Kornicker and Poore, 1996: 120*Type locality.* South of Point Hicks, Victoria, Australia, 1000 m depth. 38°21.90"S, 149°20.00"E.***Leuroleberis* Kornicker, 1981***Leuroleberis* Kornicker, 1981b: 91*Type species.* *Leuroleberis sharpei* Kornicker, 1981 by original designation.***Leuroleberis mackenziei* Kornicker, 1981***Cycloleberis bradyi* Poulsen, 1965: 268; [part; 2 juvenile females only]*Leuroleberis mackenziei* Kornicker, 1981b: 115*Type locality.* Moreton Bay, Brisbane, Australia, 22 m depth.***Leuroleberis orbicularis* (Brady, 1897)***Cyclasterope orbicularis* Brady 1897: 87.—Müller 1906: 33*Cycloleberis orbicularis*.—Skogsberg 1920: 442*Cycloleberis* species indeterminate.—Kornicker, 1975: 570*Leuroleberis orbicularis*.—Kornicker, 1981b: 92*Type locality.* Valparaiso, Chile.*Remarks.* This species is known only from juveniles of uncertain sex. It is considered to be inadequately described Kornicker (1975: 567), and may be conspecific with *Leuroleberis poulseni* (Kornicker, 1981b: 92). The holotype location is unknown.***Leuroleberis poulseni* (Mogilevsky and Ramirez, 1970)***Cycloleberis poulseni* Mogilevsky and Ramirez, 1970: 461*Leuroleberis poulseni*.—Kornicker, 1981b: 95*Type locality.* Mar del Plata, Argentina.*Remarks.* No holotype was designated. This species is possibly conspecific with *Leuroleberis orbicularis* (Kornicker, 1981b: 97).***Leuroleberis sharpei* Kornicker, 1981***Leuroleberis sharpei* Kornicker, 1981b: 97*Type locality.* Monterey Bay, California, Moss Landing Marine Laboratory, off South Jetty, 37 m depth.***Leuroleberis surugaensis* Hiruta, 1982***Leuroleberis surugaensis* Hiruta, 1982: 11

Type locality. Suruga Bay, Japan, 108-115 m depth. 35°03.3'N 138°50.0'E.

***Leuroleberis zealandica* (Baird, 1850)**

Cypridina zealandica Baird, 1850: 102
Cyclasterope zealandica.—Brady, 1898: 433
Cyclasterope tenera Brady, 1898: 433
Cyclasterope ovulum Brady, 1898: 432
Cycloleberis zealandica.—Skogsberg, 1920: 442
Cycloleberis tenera.—Skogsberg, 1920: 442
Cycloleberis ovulum.—Skogsberg, 1920: 442
Cycloleberis bradyi Poulsen, 1965: 268 [holotype only]
Azyogocypridina zealandica.—Eagar, 1971: 60 [part, only *Cypridina zealandica*]
Leuroleberis zealandica.—Kornicker, 1981b: 95

Type locality. New Zealand; exact locality unknown (Kornicker, 1975: 569).

Taxonomic decision for synonymy. Kornicker (1981b) [most recent list].

Remarks. There are two syntypes. The species is considered to be inadequately described (Kornicker, 1975: 567).

***Macroasteropteron* Kornicker, 1994**

Macroasteropteron Kornicker, 1994: 183

Type species. *Macroasteropteron mindax* Kornicker, 1994 by original designation.

***Macroasteropteron mindax* Kornicker, 1994**

Macroasteropteron mindax Kornicker, 1994: 184

Type locality. New South Wales, Australia, 996 m depth. 34°52.72'S, 151°15.04'E.

Remarks. The species is based on a unique specimen.

***Microasteropteron* Poulsen, 1965**

Microasteropteron Poulsen, 1965: 216

Type species. *Microasteropteron parvum* Poulsen, 1965 by monotypy.

***Microasteropteron bacescui* Kornicker, 1981**

Microasteropteron bacescui Kornicker, 1981b: 347

Type locality. Bab el Mandeb (strait between Red Sea and Gulf of Aden), 24 m depth.

***Microasteropteron parvum* Poulsen, 1965**

Microasteropteron parvum Poulsen, 1965: 217

Type locality. Koh Kam, Thailand, 18 m depth.

***Microasteropteron youngi* Kornicker, 1976**

Microasteropteron youngi Kornicker, 1976a: 15

Type locality. Between reef and Paiko Peninsula, Maunalua Bay, Oahu, Hawaii, 0.5 m depth.

***Monasterope* Kornicker, 1991**

Monasterope Kornicker, 1991a: 124

Type species. *Monasterope bex* Kornicker, 1991, by original designation.

***Monasterope bex* Kornicker, 1991**

Parasterope sp. Kornicker, 1987: 217

Monasterope bex Kornicker, 1991a: 125

Type locality. Enewetak Lagoon, Enewetak Atoll.

***Omegasterope* Kornicker, 1981**

Omegasterope Kornicker, 1981b: 343

Type species. *Asteropteron upsilon* Kornicker and Caraion, 1974 by original designation.

***Omegasterope upsilon* (Kornicker and Caraion, 1974)**

Asteropteron upsilon Kornicker and Caraion 1974: 59 [part]

Omegasterope upsilon.—Kornicker and Caraion, 1974: 343

Type locality. Off coast of Mauritania, 82 m depth. 20°17'03"N, 17°36'06"W.

***Parasterope* Kornicker, 1975**

Parasterope Poulsen, 1965: p [nomen nudum; without type species; 11 included species, *Parasterope obesa*, *P. pectinata*, *P. muelleri*, *P. skogsbergii* now included in *Skogsbergiella*, *P. longungues*, *P. jensenii*, *P. corrugata* now included in *Postasterope*, *P. nana*, *P. ohlini*, *P. aberrata*, *P. curta* now included in *Homasterope*]

Parasterope Kornicker, 1975: 401 [designated type species]

Type species. *Asterope Müllerii* Skogsberg, 1920 by original designation, Kornicker, 1975.

***Parasterope aberrata* (Skogsberg, 1920)**

Asterope aberrata Skogsberg, 1920: 504

Parasterope aberrata.—Poulsen, 1965: 362

Type locality. Ireland, Strangford Lough.

***Parasterope alpha* Kornicker and Caraion, 1974**

Parasterope alpha Kornicker and Caraion, 1974: 11

Type locality. Mauritania, 560 m depth. 18°40'00"N, 16°41'05"W.

***Parasterope anommata* Kornicker, 1975**

Parasterope anommata Kornicker, 1975: 427

Type locality. continental slope off Point Descado, east of Argentina, 424-428 m depth.

***Parasterope antyx* Kornicker, 1989**

Parasterope antyx Kornicker, 1989: 97

Type locality. west of Bay of Biscay, 47°39'N, 8°11'48"W, 1560 m depth.

***Parasterope beta* Kornicker, 1976**

Parasterope beta Kornicker, 1976b: 11

Type locality. Luderitz Bay, South-west Africa.

***Parasterope crinita* Kornicker, 1975**

Parasterope crinita Kornicker, 1975: 421

Type locality. Subantarctic, east of South Island, New Zealand, 51 m depth.

***Parasterope extrachelata* (Kornicker, 1958)**

Asteropina extrachelata Kornicker, 1958: 241

Synasterope extrachelata.—Poulsen, 1965: 355

Parasterope extrachelata.—Kornicker, 1986: 28

Type locality. Cavelle Pond, salt water pond on South Bimini, Great Bahama Bank.

Remarks. The species is based on a unique specimen, but the holotype is not extant. Kornicker (1986: 28) described an adult male and noted that the mandible e-seta on the female is assumed but not certain.

Parasterope gamurru Syme and Poore, 2006

Parasterope gamurru Syme and Poore, 2006: 52

Type locality. Patch Reef, Lizard Island, Queensland, Australia. 14°41.038' S, 145°27.003' E, 3 m.

Parasterope kappa Kornicker, 1976

Parasterope kappa Kornicker, 1976b: 18

Type locality. Luderitz Bay, South-west Africa.

Parasterope hirutai Chavtur, 1983

Parasterope hirutai Chavtur, 1983 [reference not sighted]

Remarks. We were unable to obtain this reference hence do not have information about the type locality.

Parasterope hulingsi Baker, 1978

Cylindroleberis psitticina Darby, 1965 : 317

Parasterope hulingsi Baker, 1978: 145

Type locality. south of San Mateo Point, California, 33°17'35"S, 117°28'50"W, 13.7 m depth.

Parasterope jensei Poulsen, 1965

Parasterope jensei Poulsen, 1965: 387

Type locality. Okinose, Sagami Sea, Japan, 180 m depth.

Parasterope lagunicola Hartmann, 1984

Parasterope lagunicola Hartmann, 1984: 121

Type locality. Kalkalgen vom Anleger der Kia Ora Village Lagune.

Parasterope longiseta (Skogsberg, 1920)

Asterope Müller var. longiseta Skogsberg, 1920 : 467

Parasterope muelleri longiseta.—Poulsen, 1965: 469

Parasterope longiseta.—Kornicker, 1975: 405

Type locality. Falkland Islands, Port William, 71 m depth.

Parasterope longungues Poulsen, 1965

Parasterope longungues Poulsen, 1965: 382

Type locality. Off Frederickssted, St. Croix, Virgin Islands, 900 m depth.

Remarks. The species is based on a unique specimen.

Parasterope lux Kornicker, 1996

Parasterope lux Kornicker, in Kornicker and Poore, 1996: 110

Type locality. South of Point Hicks, Victoria, Australia, 1000 m depth. 38°21.90'S, 149°20.00'E.

Remarks. The species is based on a unique specimen.

Parasterope maddocksa Kornicker & Thomassin, 1998

Parasterope maddocksa Kornicker and Thomassin, 1998: 117

Type locality. Tulear Reef Complex, Southwest Madagascar, Grand Recif, 24 m depth.

Parasterope mauritania Kornicker and Caraion, 1974

Parasterope mauritania Kornicker and Caraion, 1974: 18

Type locality. Mauritania, 52 m depth. 20°10'05"N, 17°32'09"W.

Parasterope mckenziei Kornicker, 1970

Parasterope mckenziei Kornicker, 1970: 26

Type locality. southwest of Botic Island, Salcedo, Samar Province, Philippines, 1 m depth. 11°05'N, 125°41'E.

Parasterope micrommata Kornicker, 1975

Parasterope micrommata Kornicker, 1975: 419

Type locality. Subantarctic, near Macquarie Island, 112 m depth.

Parasterope muelleri (Skogsberg, 1920)

Cypridina teres Norman, 1861

Asterope teres.—Müller 1912: 46 [part]

Asterope Müller Skogsberg 1920: 483

Asteropina muelleri.—Kornicker, 1958: 239

Parasterope muelleri.—Poulsen, 1965: 370

Type locality. English Channel, off Salcombe, coast of England.

Taxonomic decision for synonymy. Kornicker and Iliffe (2000) [most recent list].

Remarks. Kornicker (1974b) discussed the existing specimen descriptions. A supplementary description of the female was provided by Kornicker and Caraion (1974: 8). Another supplementary description of an adult female was provided by Kornicker, (1986: 15), who also noted that this species is probably a species complex. Kornicker and Iliffe (2000) described an additional specimen and discussed the Eastern and Western Atlantic specimens.

Parasterope nana Poulsen, 1965

Parasterope nana Poulsen, 1965: 396

Type locality. Koh Kam, Thailand, 18 m depth.

Parasterope obesa Poulsen, 1965

Parasterope obesa Poulsen, 1965: 364

Type locality. Misaki, Japan, shallow water.

Remarks. The species is based on a unique specimen.

Parasterope ohlini (Skogsberg, 1920)

Asterope Ohlini Skogsberg, 1920: 467

Parasterope ohlini.—Poulsen 1965: 363

Parasterope lowryi Kornicker, 1971: 197

Type locality. South Georgia, off Grytviken, Antarctic, 24-52 m depth. 54°22'S, 36°27'W.

Taxonomic decision for synonymy. Kornicker (1975: 402).

Parasterope pacifica Kornicker and Harrison-Nelson, 2005

Parasterope pacifica Kornicker and Harrison-Nelson, 2005: 325

Type locality. Johnston Atoll, north of Akau Island, 2-3.5 m depth. 16°45'53.3"N, 169°31'2.4"W

Parasterope pectinata Poulsen, 1965

Parasterope pectinata Poulsen, 1965: 367

Type locality. Three Kings Island, Northern New Zealand, 100 m depth.

Remarks. The species is based on one adult female and one juvenile.

Parasterope physinx Kornicker, 1996

Parasterope physinx Kornicker, in Kornicker and Poore, 1996: 112

Type locality. South of Point Hicks, Victoria, Australia, 400 m depth. 38°17.70'S, 149°11.30'E.

***Parasterope pollex* Kornicker, 1967**

Parasterope pollex Bowman and Kornicker, 1967: 9
Parasterope sp. indet. Kornicker, 1977: 794

Type locality. Hadley Harbor, Massachusetts.

Remarks. A supplementary description of the adult female was provided by Kornicker (1986).

***Parasterope prolixa* Kornicker, 1975**

Parasterope prolixa Kornicker, 1975: 436

Type locality. Subantarctic, Fortesque Bay, Strait of Magellan, 21 m depth.

***Parasterope pseudoquadrata* (Hartmann 1965)**

Cylindroleberis pseudoquadrata Hartmann, in Hartmann-Schröder and Hartmann, 1965: 316

Parasterope pseudoquadrata.—Kornicker, 1975: 415

Type locality. Bahia Ingles, North Chile, 12 m depth, 41°48'S, 73°53'W.

Remarks. A supplementary description was provided by Kornicker (1975).

***Parasterope quadrata* (Brady 1898)**

Asterope quadrata Brady, 1898: 432
Not Synasterope quadrata.—Poulsen, 1965: 406
Synasterope quadrata.—Eagar, 1971: 61 [part]
Parasterope quadrata.—Kornicker, 1975: 409

Type locality. Lyttelton Harbor, New Zealand, 2-10 m depth.

Remarks. No holotype was designated for this species.

***Parasterope quadrata?* (Brady 1898)**

Asterope australis Brady, 1898: 431
Parasterope australis.—Eagar, 1971: 61
Parasterope quadrata?.—Kornicker, 1975: 412

Remarks. Kornicker (1975: 412) provided a discussion of the specimens.

***Parasterope sequax* Kornicker, 1996**

Parasterope sequax Kornicker, in Kornicker and Poore, 1996: 106

Type locality. South of Point Hicks, Victoria, Australia, 1500 m depth. 38°25.00'S, 149°00.00'E.

***Parasterope sohni* Kornicker and Caraion, 1974**

Parasterope sohni Kornicker and Caraion, 1974: 14

Type locality. West Africa, 534 m depth. 21°47'00"N, 17°36'06"W.

***Parasterope styx* Kornicker, 1975**

Parasterope styx Kornicker, 1975: 432

Type locality. South Pacific, West of Chile, 4303m. 37°57'S, 75°08'W.

Remarks. A supplementary description of the adult female and a description of the adult male was provided by Kornicker (1989).

***Parasterope thrix* Kornicker and Caraion, 1974**

Parasterope thrix Kornicker and Caraion, 1974: 21

Type locality. Mauritania, 270 m depth. 20°50'05"N, 17°39'00"W.

Remarks. The species is based on a unique specimen.

***Parasterope whatleyi* Kornicker, 1996**

Parasterope whatleyi Kornicker, in Kornicker and Poore, 1996: 115

Type locality. New South Wales, Australia, 996 m depth. 34°52.72'S, 151°15.04'E

***Parasterope zamboangae* Kornicker, 1970**

Parasterope zamboangae Kornicker, 1970: 23

Type locality. Sacol Island, Zamboanga Province, Philippines, 1-2 m depth.

Remarks. The species is based on a unique specimen.

***Parasterope zeta* Kornicker, 1986**

Parasterope zeta Kornicker, 1986: 47

Type locality. Gulf of Mexico, off Galveston, Texas, USA, 58 m depth.

***Polyleberis* Kornicker 1974**

Polyleberis Kornicker, 1974b: 48

Type species. *Polyleberis mackenziei* Kornicker, 1974, by original designation.

***Polyleberis mackenziei* Kornicker, 1974**

Polyleberis mackenziei Kornicker, 1974b: 50

Type locality. Gulf of Naples, Italy.

***Postasterope* Kornicker, 1986**

Postasterope Kornicker, 1986: 96

Type species. *Postasterope abaco* Kornicker, 1986, by original designation.

***Postasterope abaco* Kornicker, 1986**

Postasterope abaco Kornicker, 1986: 105

Type locality. Abaco Island, Bahamas.

Remarks. The species is known from several adult males but no females.

***Postasterope barensi* (Baker, 1978)**

Parasterope barensi Baker, 1978: 139
Postasterope barensi.—Kornicker, 1986: 96

Type locality. south of Santa Barbara, California, USA, 12.8 m depth.

***Postasterope corrugata* (Poulsen, 1965)**

Parasterope corrugata Poulsen 1965: 391
Postasterope corrugata.—Kornicker, 1986: 98

Type locality. Virgin Islands, West Indies.

Remarks. The species is known from one adult male, but no females.

***Postasterope messingi* Kornicker, 1986**

Postasterope messingi Kornicker, 1986: 99

Type locality. Key Biscayne side of Bear Cut, Florida Keys, 2-4 m depth.

***Prionotoleberis* Kornicker, 1974**

Prionotoleberis (Kornicker, 1974b: 43

Type species. *Prionotoleberis gyion* Kornicker, 1974, by original designation.

Remarks. An emended diagnosis was provided by Kornicker (1992: 190).

***Prionotoleberis abyssicola* (Sars, 1870)**

Asterope abyssicola Sars, 1870: 170

Prionotoleberis abyssicola.—Kornicker, 1991b: 21

Type locality. Lofoten Islands between Finmark and Beeren Island, 290-345 m depth.

***Prionotoleberis gyion* Kornicker, 1974**

Prionotoleberis gyion Kornicker, 1974b: 43

Type locality. Gulf of Naples, Italy, near Benta Palumma, 55 m depth.

***Prionotoleberis lux* Kornicker, 1992**

Prionotoleberis lux Kornicker, 1992: 190

Type locality. near Mayotte, Indian Ocean, 300-350 m depth.

***Prionotoleberis norvegica* (Sars, 1869)**

Asterope norvegica Sars, 1869: 357

Prionotoleberis norvegica.—Kornicker, 1991b: 22

Type locality. vicinity of Sweden and Norway, 70-160 m depth.

***Prionotoleberis pax* Kornicker and Caraion, 1974**

Prionotoleberis pax Kornicker and Caraion, 1974: 37

Type locality. Eastern Atlantic Ocean off Mauritania, 270 m depth. 20°50'05" N, 17°39'00" W.

***Prionotoleberis rex* Kornicker, 1989**

Prionotoleberis rex Kornicker, 1989: 120

Type locality. Bay of Biscay, 1845-1913 m depth.

***Prionotoleberis salomani* Kornicker, 1986**

Prionotoleberis salomani Kornicker, 1986: 89

Type locality. Panama City, Florida, 9.1 m depth.

***Prionotoleberis styx* Kornicker, 1991**

Prionotoleberis styx Kornicker, 1991b: 22

Type locality. Eastern Pacific Ocean, hydrothermal vent, 2592 m depth. 20°51'00" N, 109°04'00" W.

***Pteromeniscus* Kornicker, 1981**

Pteromeniscus Kornicker, 1981b: 333

Type species. *Pteromeniscus intesi* Kornicker, 1981, by original designation.

***Pteromeniscus intesi* Kornicker, 1981**

Pteromeniscus intesi Kornicker, 1981b: 334

Type locality. Mauritanian, continental shelf, 699-270 m depth.

Remarks. The species is known only from an adult male and juvenile female.

***Skogsbergiella* Kornicker, 1975**

Skogsbergiella Kornicker, 1975: 465

Philippiella Poulsen, 1965: 344 [part] [nomen nudum, without type species; 2 included species, *Asterope quinquesetae* now included in *Archasterope*, and *Asterope spinifera* now in *Skogsbergiella*]

Type species. *Asterope spinifera* Skogsberg, 1920 by original designation.

Remarks. An emended diagnosis was provided by Kornicker (in Kornicker and Poore, 1996: 161).

***Skogsbergiella inermis* Chavtur, 1983**

Skogsbergiella inermis Chavtur, 1983 [reference not sighted]

Remarks. We were unable to obtain this reference do not have information about the type locality.

***Skogsbergiella macrothrix* Kornicker, 1975**

Skogsbergiella macrothrix Kornicker, 1975: 477

Type locality. Scotia Sea, Subantarctic, Magellanic subregion, 578 m depth.

***Skogsbergiella pax* Kornicker, 1975**

Skogsbergiella pax Kornicker, 1975: 495

Type locality. Argentina shelf and slope, 40-45°S, 1475 m depth.

***Skogsbergiella plocus* Kornicker, 1975**

Skogsbergiella plocus Kornicker, 1975: 489

Type locality. off Golfo de Penas, west of Chile, 2657 m depth.

***Skogsbergiella scotti* Kornicker, 1975**

Skogsbergiella scotti Kornicker, 1975: 471

Type locality. Weddell Sea, Antarctic, Scotia subregion, Arthur Harbor, Anvers Island, Palmer Station, 6 m depth.

***Skogsbergiella senex* Kornicker, 1996**

Skogsbergiella senex Kornicker, in Kornicker and Poore, 1996: 161

Type locality. South of Point Hicks, Victoria, Australia, 1850 m depth. 38°25.90'S, 148°58.60'E.

***Skogsbergiella skogsbergi* (Kornicker, 1971)**

Philippiella skogsbergi Kornicker, 1971: 211

Skogsbergiella skogsbergi.—Kornicker, 1975: 471

Type locality. Vicinity of Greenwich Island, South Shetland Islands, 274 m depth.

***Skogsbergiella spinifera* (Skogsberg, 1920)**

Asterope spinifera Skogsberg, 1920: 466

Philippiella spinifera.—Poulsen, 1965: 344

Skogsbergiella spinifera.—Kornicker, 1975: 467

Type locality. South Georgia, off mouth of Cumberland Bay. 54°11'S, 54°18'W.

***Synasterope* Kornicker, 1975**

Synasterope Poulsen, 1965: 401 [nomen nudum; without type species; 10 included species, *Synasterope bassana*, *S. quadrata*, *S. quatrisetosa*, *S. knudseni*, *S. oculata*, *S. longiseta*, *S. implumis*, *S. serrata*, *S. setispara*, *S. extrachelata* now in *Parasterope*]

Synasterope Kornicker, 1975: 440 [designated type species]

Type species. *Synasterope implumis* Poulsen, 1965, by original designation.

***Synasterope arnaudi* Kornicker, 1975**

Synasterope arnaudi Kornicker, 1975: 463

Type locality. Antarctic, Kerguelen Islands, 50 m depth.

***Synasterope bassana* Poulsen, 1965**

Synasterope bassana Poulsen, 1965: 402

Type locality. Bass Strait, southeast Australia, 180 m depth. 38°12'S, 149°40'E.

Remarks. No holotype not designated for this species.

Synasterope bathyalis* Chavtur, 1983Synasterope bathyalis* Chavtur, 1983 [reference not sighted]

Remarks. We were unable to obtain this reference and do not have information about the type locality.

Synasterope bensoni* Kornicker and Caraion, 1974Synasterope bensoni* Kornicker and Caraion, 1974: 23

Type locality. Mauritania, 270 m depth. 20°50'05" N, 17°39'00" W.

Synasterope brachythrix* Kornicker, 1975Synasterope brachythrix* Kornicker, 1975: 461

Type locality. Subantarctic, Magellanic subregion, in channel off Puerto Bheno, 214 m depth.

?*Synasterope brevisetae* Hartmann, 1985*Synasterope brevisetae* Hartmann, 1985: 98

Type locality. Indian Ocean.

Remarks. This reference was not sighted. This information is from Kornicker (1992: 194).

Synasterope browni* Kornicker and Iliffe, 2000Synasterope browni* Kornicker and Iliffe, 2000: 72

Type locality. Exuma Sound, Bahamas, 96 m depth.

Remarks. A description of an adult male is provided by Kornicker and Harrison-Nelson (2002: 85).

Synasterope calix* Kornicker, 1992Synasterope calix* Kornicker, 1992: 194

Type locality. Mozambique Channel, Zelee Bank, 18-24 m depth.

Remarks. Supplementary descriptions of an adult female and adult male were provided by Kornicker and Thomassin (1998).

Synasterope cushmani* Kornicker, 1974Synasterope cushmani* Kornicker, 1974a: 11

Type locality. Cape Cod Bay, Massachusetts, 36 m depth.

***Synasterope dimorpha* (Hartmann, 1965)**

Cylindroleberis dimorpha Hartmann, in Hartmann-Schröder and Hartmann, 1965: 321
Synasterope dimorpha.—Kornicker, 1975: 441

Type locality. near Cabo Quedal, Chile, 136 m depth.

Synasterope duplex* Kornicker, 1975Synasterope duplex* Kornicker, 1975: 448

Type locality. Weddell Sea, Antarctica, 3035 m depth.

Synasterope empoulseni* Kornicker, 1975Synasterope empoulseni* Kornicker, 1975: 447

Type locality. Halfmoon Bay, Stewart Island, New Zealand, 10-15 m depth.

Synasterope helix* Kornicker, 1989Synasterope helix* Kornicker, 1989: 111

Type locality. west of Bay of Biscay, 2165 m depth. 47°33'42" N, 8°39'54" W.

Synasterope hirpex* Kornicker, 1989Synasterope hirpex* Kornicker, 1989: 105

Type locality. west of Bay of Biscay, 4706 m depth. 46°31'12" N, 10°23'48" W.

Synasterope implumis* Poulsen, 1965Synasterope implumis* Poulsen, 1965: 421

Type locality. Long Point Bay, Virgin Islands, West Indies, shallow water.

Synasterope index* Kornicker, 1989Synasterope index* Kornicker, 1989: 115

Type locality. west of Bay of Biscay, 2906m. 47°30'48" N, 9°07'36" W.

Remarks. The species is based on a juvenile male only.

Synasterope longiseta* Poulsen, 1965Synasterope longiseta* Poulsen, 1965: 417

Type locality. Virgin Islands, West Indies, depth unknown.

Remarks. The species is based on a unique specimen.

Synasterope mystax* Kornicker, 1975Synasterope mystax* Kornicker, 1975: 456

Type locality. Scotia Sea, Antarctic, 2747 m depth.

Synasterope oculata* (Brady, 1902)Asterope oculata* Brady, 1902: 179*Synasterope oculata*.—Poulsen, 1965: 401

Type locality. Trincomalee, Sri Lanka.

Remarks. Kornicker (1992: 222) discussed the specimens used by Brady (1902), Skogsberg (1920) and Poulsen (1965).

Synasterope phalanx* Kornicker, 1989Synasterope phalanx* Kornicker, 1989: 102

Type locality. west of Bay of Biscay, 2147 m depth. 47°32'42" N, 8°36'30" W.

Remarks. The species is known only from adult males.

Synasterope polythrix* Kornicker, 1975Synasterope polythrix* Kornicker, 1975: 454

Type locality. Drake Passage, Antarctica, 4004 m depth.

Synasterope psitticina* (Darby, 1965)Cylindroleberis psitticina* Darby, 1965: 31 [part]*Synasterope psitticina*.—Kornicker, 1974a: 18

Type locality. Continental shelf off Sapelo Island, Georgia, exact depth unknown but between 13-123m.

Remarks. A supplementary description of the adult female holotype was provided by Kornicker (1974a).

Synasterope quadrata* (Brady, 1898)Synasterope quadrata*.—Poulsen, 1965: 406*Not Asterope quadrata* Brady, 1898: 432*Not Parasterope quadrata* Kornicker, 1975: 409

Taxonomic decision for synonymy. Kornicker (1975: 409).

Type locality. No type specimen. Specimens from New Zealand.

Remarks. Poulsen (1965) described an adult female based on new material, which he believed was conspecific with *Asterope quadrata* Brady, 1898. He then placed the species in *Synasterope*. However, Kornicker (1975) re-examined Brady's material and placed the species in *Parasterope*. He excluded

Poulsen's specimens from the synonymy. Hence, we believe that Poulsen's specimens – if found to represent a new species – should be re-named as a new species, not as *quadrata*.

***Synasterope quatriisetosa* Poulsen, 1965**

Synasterope quatriisetosa Poulsen, 1965: 410

Type locality. Apia Harbour, Samoa Island, surface.

Remarks. This species is described only from one adult male.

***Synasterope serrata* Poulsen, 1965**

Synasterope serrata Poulsen, 1965: 427

Type locality. Northeast of Hans Lollik, St. Thomas, Virgin Islands, West Indies, 50 m depth.

Remarks. The species is based on a unique specimen.

***Synasterope setispara* (Kornicker, 1958)**

Asteropina setispara Kornicker, 1958: 239

Synasterope setispara.—Poulsen, 1965: 396

Type locality. Bimini area, Great Bahama Bank.

Remarks. The holotype is considered to be lost. A lectotype was designated by Kornicker (1986).

***Synasterope solox* Kornicker, 1996**

Synasterope solox Kornicker, in Kornicker and Poore, 1996: 166

Type locality. off Eden, New South Wales, Australia, 220 m depth. 36°57.40'S, 150°18.80'E.

***Synasterope williamsae* Kornicker, 1986**

Synasterope williamsae Kornicker, 1986: 80

Type locality. Gulf of Mexico, off Galveston, Texas, USA, 55 m depth.

***Tetraleberis* Kornicker, 1981**

Tetraleberis Kornicker, 1981b: 134

Type species. *Asterope brevis* Müller, 1890, by original designation.

***Tetraleberis brevis* (Müller, 1890)**

Asterope brevis Müller, 1890: 239

Cyclasterope brevis.—Müller, 1906: 33

Cylindroleberis brevis.—Kajiyama, 1912: 617

Cycloleberis brevis.—Skogsberg, 1920: 442

Tetraleberis brevis.—Kornicker, 1981b: 135

Type locality. Coast of Japan, near Enoshima, 22 m depth.

Remarks. There are 16 syntypes.

***Tetraleberis maddockiae* Kornicker, 1981**

Tetraleberis maddockiae Kornicker, 1981b: 146

Type locality. Madagascar, Grand Recif, Grand Vasque, 18 m depth.

***Tetraleberis pix* Kornicker, 1996**

Tetraleberis pix Kornicker, 1996: 79

Type locality. Calliope River and Auckland Creek area, near Gladstone, Queensland, Australia.

***Tetraleberis similis* (Brady, 1902)**

Cyclasterope similis Brady, 1902: 183

Cycloleberis similis.—Poulsen, 1965: 243

Tetraleberis similis.—Kornicker, 1981b: 143

Type locality. Java Sound, 14.6 m depth.

***Tetraleberis tanzania* Kornicker, 1981**

Tetraleberis tanzania Kornicker, 1981b: 153

Type locality. Off Tanzania. no station data.

***Tetraleberis triplex* Kornicker, 1996**

Tetraleberis triplex Kornicker, 1996: 89

Type locality. Lizard Island, Australia, 4 m depth.

***Xandarasterope* Kornicker, 1996**

Xandarasterope Kornicker, in Kornicker and Poore, 1996: 126

Type species. *Xandarasterope storthynx* Kornicker, 1996, by original designation.

***Xandarasterope storthynx* Kornicker, 1996**

Xandarasterope storthynx Kornicker, in Kornicker and Poore, 1996: 126

Type locality. South of Point Hicks, Victoria, Australia, 2900 m depth. 38°40.29'S, 149°18.06"E.

***Xandarasterope trux* Kornicker, 1996**

Xandarasterope trux Kornicker, in Kornicker and Poore, 1996: 131

Type locality. South of Point Hicks, Victoria, Australia, 1840 m depth. 38°29.33'S, 149°19.98"E.

***Xenoleberis* Kornicker, 1994**

Xenoleberis Kornicker, 1994: 177

Type species. *Xenoleberis bex*, Kornicker, 1994, by original designation.

***Xenoleberis bex* Kornicker, 1994**

Xenoleberis bex Kornicker, 1994: 178

Type locality. South of Point Hicks, Victoria, Australia, 600 m depth. 38°19.10'S, 149°14.30"E.

Remarks. The species is based on a unique specimen.

***Xenoleberis californica* (Baker, 1979)**

Bathyleberis californica Baker, 1979: 287

Xenoleberis californica.—Kornicker, 1994: 177

Type locality. north of Point Conception, California, 50 m depth.

***Xenoleberis yamadai* (Hiruta, 1979)**

Xenoleberis yamadai Hiruta, 1979: 99

Bathyleberis yamadai.—Kornicker, 1994: 177

Type locality. Oshoro, Japan, 3-5 m depth.

References

- Baird, W. 1850. Description of several new species of Entomostraca. *Proceedings of the Zoological Society of London* 18: 254-257.
- Baird, W. 1860. Note upon the genus *Cypridina*, Milne-Edwards, with a description of some new species. *Proceedings of the Zoological Society London* 28: 199-202.
- Baker, J.H. 1978. Two new species of *Parasterope* (Myodocopina, Ostracoda) from southern California. *Crustaceana* 35: 139-151.
- Baker, J.H. 1979. Three new species of *Bathyleberis* (Ostracoda, Myodocopina) from southern California, USA. *Crustaceana* 36: 287-301.
- Bowman, T.E., and Kornicker, L.S. 1967. Two new crustaceans: the parasitic copepod *Sphaeronellopsis monothrix* (Choniostomatidae) and its myodocopid ostracod host *Parasterope pollex*

- (Cylindroleberididae) from the Southern New England coast. *Proceedings of the United States National Museum* 123: 29.
- Brady, G.S. 1868. A synopsis of the recent British Ostracoda. *The Intellectual Observer* 12: 110-130.
- Brady, G.S. 1871. A review of the Cypridinidae of the European Seas, with description of a new species. *Proceedings of the Zoological Society of London* 19: 289-296.
- Brady, G.S. 1897. A supplementary report on the Crustaceans of the group Myodocopa obtained during the Challenger Expedition, with notes on other new or imperfectly known species. *Transactions of the Zoological Society of London* 14: 85-100.
- Brady, G.S. 1898. On new or imperfectly known species of Ostracoda, chiefly from New Zealand. *Transactions of the Zoological Society of London* 14: 489-525.
- Brady, G.S. 1902. On new or imperfectly known Ostracoda, chiefly from a collection in the Zoological Museum, Copenhagen. *Transactions of the Zoological Society of London* 16: 179-210.
- Brady, G.S., and Norman, A.M. 1896. A monograph of the marine and freshwater Ostracoda of the North Atlantic and North-Western Europe. *The Scientific Transactions of the Royal Dublin Society* 2: 621-784.
- Chavtur, V.G. 1978. *Euphilomedes nipponica* Hiruta, *Scleroconcha ochotensis* n. sp. and *Empoulsenia kurilensis* n. sp. (Ostracoda, Myodocopina) from the region of the Kurile Islands. Pp. 149-158 in: Kussakin, O.G. (ed.) *Fauna and Flora of the Shelf of the Kurile Islands (In Russian, with English abstract)*.
- Chavtur, V.G. 1983. [reference not sighted] Ostrakody Myodocopina, Cladocopina umerennykh I kholodnykh vod Severnogo polusharila. [Ostracodes (Myodocopina, Cladocopina) of Temperate and Cold Waters of the Northern Hemisphere]. *Academy of Sciences of the USSR, Far-Eastern Science Center, Institute of Marine Biology*: 132 pp.
- Claus, C. 1876. *Untersuchungen zur Erforschung der genealogischen Grundlage des Crustaceen-System*. Wien.
- Cohen, A.C., and Kornicker, L.S. 1975. Taxonomic Indexes to Ostracoda (Suborder Myodocopina) in Skogsberg (1920) and Poulsen (1962, 1965). *Smithsonian Contributions to Zoology* 204: 29.
- Darby, D. 1965. Ecology and taxonomy of Ostracoda in the vicinity of Sapelo Island, Georgia. Pp. 1-77 in: Kesling, R.V. (ed.) *Four Reports of Ostracod Investigations*.
- Eagar, S.H. 1971. A check list of the Ostracoda of New Zealand. *Journal of the Royal Society of New Zealand* 1: 53-64.
- Granata, L. 1915. Nuove Specie di Ostracodi. *Bullettino della Societa Entomologica Italiana* 46.
- Hanai, T. 1959. Studies on the Ostracoda from Japan: Historical review with bibliographic index of Japanese Ostracoda. *Journal of the Faculty of Science, The University of Tokyo* 11: 419-439.
- Hartmann, G. 1974. Zur Kenntnis des Eulitorals der afrikanischen Westküste zwischen Angola und Kap der Guten Hoffnung und der afrikanischen Ostküste von Südafrika und Moçambique unter besonderer Berücksichtigung der Polychaeten und Ostracoden, part 3: Die Ostracoden des Untersuchungsgebiets. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 69: 229-520.
- Hartmann, G. 1984. Zur Kenntnis der Ostracoden der polynesischen Inseln Huahine (Gesellschaftsinseln) und Rangiroa (Tuamotu-Inseln). *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 81: 117-169.
- Hartmann, G. 1985. [reference not sighted] Ostracoden aus der Tiefsee des Indischen Ozeans und der Iberischen See sowie von östatlantischen sublitoralen Plateaus und Kuppen. *Senckenbergiana Maritima* 17: 89-146.
- Hartmann, G. 1989. Antarctic benthic Ostracoden IV. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 86: 209-230.
- Hartmann-Schröder, G., and Hartmann, G. 1965. Zur Kenntnis des Sublitorals der chilenischen Küste unter besonderer Berücksichtigung der Polychaeten und Ostracoden. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 62: 1-384.
- Hiruta, S. 1979. A new species of *Bathyleberis* Kornicker from Hokkaido, with reference to the larval stages (Ostracoda: Myodocopina). *Journal of the Faculty of Science, Hokkaido University* 22: 99-121.
- Hiruta, S. 1982. A new species of the genus *Leuroleberis* Kornicker from Suruga Bay, Central Japan (Ostracoda: Myodocopina). *Journal of Hokkaido University of Education* 33: 11-18.
- Kajiyama, E. 1912. The Ostracoda of Misaki, Part 2. *Dobutsugaku-zadshi* 24: 609-619.
- Kempf, E.K. 1986a. *Index and bibliography of marine Ostracoda Part 1: Index A [names of genera in alphabetical order]*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. 762 pp.
- Kempf, E.K. 1986b. *Index and bibliography of marine Ostracoda Part 2: Index B [names of species in alphabetical order]*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. 712 pp.
- Kempf, E.K. 1987. *Index and bibliography of marine Ostracoda Part 3: Index C [names of taxa in alphabetical order]*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. 774 pp.
- Kempf, E.K. 1988. *Index and bibliography of marine Ostracoda Part 4: Bibliography A [references to taxonomic publications]*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. 454 pp.
- Kempf, E.K. 1994. *Index and bibliography of marine Ostracoda Part 5: Bibliography B [references to non-taxonomic publications]*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. 366 pp.
- Kempf, E.K. 1995a. *Index and bibliography of marine Ostracoda Part 6: Index A, Supplement 1*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. 244 pp.
- Kempf, E.K. 1995b. *Index and bibliography of marine Ostracoda Part 7: Index B, Supplement 1*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. 206 pp.
- Kempf, E.K. 1995c. *Index and bibliography of marine Ostracoda Part 8: Index C, Supplement 1*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. 230 pp.
- Kempf, E.K. 1996. *Index and bibliography of marine Ostracoda Part 9: Bibliography C [references to taxonomic publications]*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. 198 pp.
- Kempf, E.K. 2002. *Index and bibliography of marine Ostracoda Part 10: Bibliography D [references to non-taxonomic publications]*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. CD-ROM pp.
- Kempf, E.K. 2004. *Index and bibliography of marine Ostracoda Part 16: Index D [genera and species]*. Geologisches Institut der Universität Köln, Sonderveröffentlichungen: Cologne. CD-ROM pp.
- Kornicker, L.S. 1958. Ecology and taxonomy of recent marine Ostracodes in the Bimini Area, Great Bahama Bank. *Publications of the Institute of Marine Science* 5: 194-300.
- Kornicker, L.S. 1970. Myodocopid Ostracoda (Cypridinacea) from the Philippine Islands. *Smithsonian Contributions to Zoology* 39: 1-32.
- Kornicker, L.S. 1971. Benthic Ostracoda (Myodocopina: Cypridinacea) from the South Shetland Islands and the Palmer Archipelago, Antarctica. *Antarctic Research Series* 17: 167-216.
- Kornicker, L.S. 1974a. Ostracoda (Myodocopina) of Cape Cod Bay, Massachusetts. *Smithsonian Contributions to Zoology* 173: 1-20.
- Kornicker, L.S. 1974b. Revision of the Cypridinacea of the Gulf of Naples (Ostracoda). *Smithsonian Contributions to Zoology* 178: 1-64.
- Kornicker, L.S. 1975. Antarctic Ostracoda (Myodocopina). *Smithsonian Contributions to Zoology* 163: 1-720.
- Kornicker, L.S. 1976a. Benthic marine Cypridinacea from Hawaii (Ostracoda). *Smithsonian Contributions to Zoology* 231: 24.
- Kornicker, L.S. 1976b. Myodocopid Ostracoda from Southern Africa. *Smithsonian Contributions to Zoology* 214: 1-39.
- Kornicker, L.S. 1981a. Benthic marine Cypridinoidea from Bermuda (Ostracoda). *Smithsonian Contributions to Zoology* 331: 1-15.
- Kornicker, L.S. 1981b. Revision, distribution, ecology, and ontogeny of the ostracode subfamily Cyclasteropinae (Myodocopina: Cylindroleberididae). *Smithsonian Contributions to Zoology* 319: 1-548.
- Kornicker, L.S. 1986. Cylindroleberididae of the Western North Atlantic and the Northern Gulf of Mexico, and Zoogeography of the Myodocopina (Ostracoda). *Smithsonian Contributions to Zoology* 425: 139.
- Kornicker, L.S. 1987. Ostracoda (Myodocopina) of Enewetak Atoll. Pp. 217-219 in: Devaney, D.M., Reese, E.S., Burch, B.L., and Helfrich, P. (eds), *The Natural History of Enewetak Atoll, Volume II, Biogeography and Systematics*. U.S. Department of Energy, Office of Scientific and Technical Information, Oak Ridge, Tennessee

- Kornicker, L.S. 1988. Myodocopid Ostracoda of the Beaufort Sea, Arctic Ocean. *Smithsonian Contributions to Zoology* 456: 1-40.
- Kornicker, L.S. 1989. Bathyal and abyssal myodocopid Ostracoda of the Bay of Biscay and vicinity. *Smithsonian Contributions to Zoology* 467: 1-134.
- Kornicker, L.S. 1991a. Myodocopid Ostracoda of Enewetak and Bikini Atolls. *Smithsonian Contributions to Zoology* 505: 1-140.
- Kornicker, L.S. 1991b. Myodocopid Ostracoda of hydrothermal vents in the eastern Pacific Ocean. *Smithsonian Contributions to Zoology* 516: 46.
- Kornicker, L.S. 1992. Myodocopid Ostracoda of the Bentchedi Expedition, 1977, to the NE Mozambique Channel, Indian Ocean. *Smithsonian Contributions to Zoology* 531: 243.
- Kornicker, L.S. 1993. *Antarctic and Subantarctic Myodocopina (Ostracoda)*. Vol. 5. Koeltz Scientific Books: Koenigstein, Germany; Champaign, Ill.
- Kornicker, L.S. 1994. Ostracoda (Myodocopina) of the SE Australian continental slope, part 1. *Smithsonian Contributions to Zoology* 553: 1-200.
- Kornicker, L.S. 1996. Ostracoda (Myodocopina) from shallow waters of the Northern Territory and Queensland, Australia. *Smithsonian Contributions to Zoology* 578: 1-97.
- Kornicker, L.S., and Caraion, F.E. 1974. West African Myodocopid Ostracoda (Cylindroleberididae). *Smithsonian Contributions to Zoology* 179: 78.
- Kornicker, L.S., and Maddocks, R.F. 1977. *Cycloleberis christiei*, a new species of marine Ostracoda (Suborder Myodocopina) from Salanha Bay and Langebaan Lagoon, South Africa. *Proceedings of the Biological Society of Washington* 90: 894-914.
- Kornicker, L.S., and Poore, G.C.B. 1996. Ostracoda (Myodocopina) of the SE Australian continental slope, part 3. *Smithsonian Contributions to Zoology* 573: 1-186.
- Kornicker, L.S., and Thomassin, B.A. 1998. Ostracoda (Myodocopina) of Tulear Reef Complex, SW Madagascar. *Smithsonian Contributions to Zoology* 595: 1-134.
- Kornicker, L.S., and Iliffe, T.M. 2000. Myodocopid ostracoda from Exuma Sound, Bahamas, and from marine caves and blue holes in the Bahamas, Bermuda, and Mexico. *Smithsonian Contributions to Zoology* 606: 1-95.
- Kornicker, L.S., and Harrison-Nelson, E. 2002. Ostracoda (Myodocopa) from Bahamian blue holes. *Smithsonian Contributions to Zoology* 616: 1-99.
- Kornicker, L.S., and Harrison-Nelson, E. 2005. Ostracoda from Johnston Atoll, Pacific Ocean, and proposal of a new tribe, Bruuniellini (Myodocopina: Cylindroleberididae). *Pacific Science* 59: 323-362.
- Kornicker, L.S., and Harrison-Nelson, E. 2006. Ostracoda (Myodocopina) of Tutuila, American Samoa. *Pacific Science* 60: 243-259.
- Lalana, R., and Kornicker, L.S. 1997. *Amboleberis cubensis*, a new species of myodocopine ostracode from the vicinity of Cuba (Crustacea: Ostracoda: Cylindroleberididae). *Proceedings of the Biological Society of Washington* 110: 107-114.
- McKenzie, K.G. 1965. Myodocopid Ostracoda (Cypridinacea) from Scammon Lagoon, Baja California, Mexico, and Their Ecologic Associations. *Crustaceana* 9: 57-70.
- Milne-Edwards, H. 1840. *Histoire naturelle des Crustaces. Tome troisième*. Paris.
- Moguilevsky, A., and Ramirez, F.C. 1970. *Cycloleberis poulseni* especie nueva (Ostracoda, Asteropidae) crustaceo bentonico del area de Mar del Plata, Argentina. *Physis* 29: 461-471.
- Müller, F. 1870. Bemerkungen über Cypridina. *Jenaische Zeitschrift für Medicin und Naturwissenschaft* 5: 255-276.
- Müller, G.W. 1890. Neue Cypridiniden. *Zoologische Jahrbücher* 5: 211-252.
- Müller, G.W. 1893. Über Lebensweise und Entwicklungsgeschichte der Ostracoden. *Sitzungsberichte der Königlich Prüssischen Akademie der Wissenschaften zu Berlin* 23: 355-381.
- Müller, G.W. 1894. Die Ostracoden des Golfes von Neapel und der angrenzenden Meeres-Abschritte. *Fauna und Flora des Golfes von Neapel* 21: 404.
- Müller, G.W. 1906. Die Ostracoda der Siboga Expedition. *Siboga-Expedition* 30.
- Müller, G.W. 1908. Die Ostracoden der Deutschen Südpolar-Expedition 1901-1903. Bd. X. *Zoologie*.
- Müller, G.W. 1912. Ostracoda. *Das Tierreich* 31.
- Norman, A.M. 1861. Contributions to British Carcinology. *Annals and Magazine of Natural History*. Third Series. volume 8.
- Philippi, A. 1840. *Asterope*, ein Neues Genus der Ostracoden. *Archiv für Naturgeschichte* 6: 186-188.
- Poulsen, E.M. 1965. Ostracoda-Myodocopa part 2: Cypridiniformes-Rutidermatidae, Sarsiellidae and Asteropidae. *Dana Report* 65: 1-483.
- Sars, G.O. 1865. Oversigt af Norge marine Ostracoder. *Forhandlinger I Vidensk.-Selskabet I Christiania*. 8: 1-130.
- Sars, G.O. 1869. Undersøgelser over Christianiafjordens Dybvandsfauna. *Nyt Magazin for Naturvidenskaberne* 16: 305-362.
- Sars, G.O. 1870. Nye Dybvandsrustaceer fra Lofoten. *Forhandlinger I Vidensk.-Selskabet I Christiania*. 12 (1869): 167-174.
- Sars, G.O. 1887. Nye bidrag til kundskaben om Middelhavets Invertebratfauna: Ostracoda Mediterranea. *Archiv for Mathematik og Naturvidenskab, Christiania*. 4.
- Scott, T. 1894. Report on Entomostraca from the Gulf of Guinea, Collected by John Rattray, B.Sc. *Transactions of the Linnean Society, series 2 (Zoology)* 6: 1-161.
- Skogsberg, T. 1920. Studies on marine Ostracoda. *Zoologiska Bidrag fran Uppsala* 1: 784.
- Stebbing, T.R. 1899. The Karkinokosm, or world of Crustacea, VII. *Knowledge* 22: 29-31.
- Stebbing, T.R. 1900. On Crustacea brought by Dr. Willey from the South Seas. *Zoological Results* 5: 605-690.
- Strand, E. 1928. Miscellanea Nomenclatorica Zoologica et Palaeontologica (1-2). *Archiv für Naturgeschichte* 92 (1926): 30-75.
- Sylvester-Bradley, P.C. 1961. Myodocopida. Pp. 387-406 in: Moore, R.C. (ed.) *Treatise on Invertebrate Paleontology, Part Q*. Geological Society of America and University of Kansas Press: Kansas.
- Syme, A.E., and Poore, G.C.B. 2006. Three new ostracod species from coastal Australian waters (Crustacea: Ostracoda: Myodocopa: Cylindroleberididae). *Zootaxa* 1305: 51-67.
- Tressler, W.L. 1949. Marine Ostracoda from Tortugas, Florida. *Journal of the Washington Academy of Sciences* 39: 335-343.
- Tuel, M.D., Wilderman, M.P., Trapani, J.E., and Hayes, P.E. 1976. A biological survey of Pillar Point Harbor, El Granada, California. Redwood City, California: Marine Ecological Institute. 254.

List of genera, species and synonyms

(alphabetical by specific epithet; where applicable, current name in square brackets)

- abaco*, *Postasterope*
- aberrata*, *Asterope* [*Parasterope*]
- aberrata*, *Parasterope*
- abyssicola*, *Asterope* [*Prionotoleberis*]
- abyssicola*, *Prionotoleberis*
- Actinoseta*
- Agassizi*, *Asteropteron* [*Asteropella*]
- agassizi*, *Cyclasterope* [*Asteropella*]
- agassizii*, *Asteropella*
- Agassizii*, *Cylindroleberis* [*Asteropella*]
- Agassizii*, *Cypridina* [*Asteropella*]
- albomaculata*, ?*Cyclasterope*
- albo-maculata*, *Cypridina* [?*Cyclasterope albomaculata*]
- albomaculata*, *Cypridina* [?*Cyclasterope*]
- alpha*, *Bruuniella*
- alpha*, *Parasterope*
- Alphaleberis*
- alphathrix*, *Alphaleberis*
- altrix*, *Archasterope*
- Amboleberis*
- americana*, *Amboleberis*
- americana*, *Asterope* [*Amboleberis*]
- americana*, *Cyclasterope* [*Amboleberis*]
- americana*, *Cycloleberis* [*Amboleberis*]
- americana*, *Cylindroleberis* [*Amboleberis*]
- anommata*, *Parasterope*
- antarctica*, *Archasterope*
- antarctica*, *Empoulsenia* [*Archasterope*]
- antyx*, *Amboleberis*
- antyx*, *Parasterope*
- apex*, *Archasterope*
- Archasterope*
- arnaudi*, *Synasterope*
- arthuri*, *Asterope* [*Cyclasterope*]
- arthuri*, *Cyclasterope*
- Asterope* [*Cylindroleberis*]
- Asteropella*
- Asteropina* [*Cylindroleberis*]
- Asteropteron*
- Asteropterygion*
- australis*, *Asterope* [*Parasterope quadrata*?]
- australis*, *Parasterope* [*Parasterope quadrata*?]
- babax*, *Bathyleberis*
- bacescui*, *Cylindroleberis*
- bacescui*, *Microasteropteron*
- barensi*, *Parasterope* [*Postasterope*]
- barensi*, *Postasterope*
- bassana*, *Synasterope*
- bathyalis*, *Synasterope*
- Bathyleberis*
- bensoni*, *Synasterope*
- beta*, *Bruuniella*
- beta*, *Parasterope*
- bex*, *Monasterope*
- bex*, *Xenoleberis*
- biminiensis*, *Cyclasterope* [*Amboleberis americana*]
- biminiensis*, *Cycloleberis* [*Amboleberis americana*]
- bisetosa*, *Cyclasterope*
- bisetosa*, *Diasterope*
- brachythrix*, *Synasterope*
- bradyi*, *Cycloleberis* [*Leuroleberis zealandica*]
- breviata*, *Bruuniella*
- brevis*, *Asterope* [*Tetraleberis*]
- brevis*, *Cyclasterope* [*Tetraleberis*]
- brevis*, *Cycloleberis* [*Tetraleberis*]
- brevis*, *Cylindroleberis* [*Tetraleberis*]
- brevis*, *Tetraleberis*
- brevisetae*, ?*Synasterope*
- browni*, *Synasterope*
- Bruuniella*
- bulla*, *Archasterope*
- californica*, *Bathyleberis* [*Xenoleberis*]
- californica*, *Xenoleberis*
- calix*, *Synasterope*
- canina*, *Diasterope*
- chelisparsa*, *Actinoseta*
- christiei*, *Cycloleberis*
- climax*, *Asteropterygion*
- corrugata*, *Parasterope* [*Postasterope*]
- corrugata*, *Postasterope*
- crinita*, *Parasterope*
- cubensis*, *Amboleberis*
- curta*, *Asterope* [*Homasterope*]
- curta*, *Homasterope*
- curta*, *Parasterope* [*Homasterope*]
- cushmani*, *Synasterope*
- Cyclasterope*
- Cycloleberis*
- Cylindroleberis*
- dayi*, *Asteropterygion*
- dentata*, *Archasterope*
- Diasterope*
- dimorpha*, *Cylindroleberis* [*Synasterope*]
- dimorpha*, *Synasterope*
- Dolasterope*
- Domromeus*
- duplex*, *Synasterope*
- efficax*, *Archasterope*
- empoulseni*, *Synasterope*
- Empoulsenia* [*Archasterope*]
- extrachelata*, *Asteropina* [*Parasterope*]
- extrachelata*, *Parasterope*
- fascigera*, *Asterope* [*Cyclasterope*]
- fascigera*, *Cyclasterope*
- fascigera*, *Cycloleberis* [*Cyclasterope*]
- fascigera*, *Cylindroleberis* [*Cyclasterope*]
- fusca*, *Asterope* [*Asteropteron fuscum*]
- fusca*, *Cyclasterope* [*Asteropteron fuscum*]
- fusca*, *Cylindroleberis* [*Asteropteron fuscum*]
- fuscum*, *Asteropteron*
- galatheae*, *Cycloleberis*
- gamurru*, *Cylindroleberis*
- garthi*, *Bathyleberis*
- glacialis*, *Asterope* [*Homasterope*]
- glacialis*, *Homasterope*
- grimaldi* var. *vicina*, *Asterope* [*Cylindroleberis*]
- grimaldi* var. *vicina*, *Asteropina* [*Cylindroleberis*]
- grimaldi* var. *vicina*, *Cylindroleberis*
- grimaldi*, *Asterope* [*Cylindroleberis*]
- grimaldi*, *Asteropina* [*Cylindroleberis*]
- grimaldi*, *Cylindroleberis*
- grisea*, *Asterope* [*Diasterope*]
- grisea*, *Diasterope*
- grossmani*, *Bathyleberis*
- gyion*, *Prionotoleberis*
- hancocki*, *Bathyleberis*
- hazeli*, *Bruuniella*
- helix*, *Synasterope*
- hendersoni*, *Cyclasterope*
- heptathrix*, *Domromeus*
- Heptonema*
- hilgendorfii*, *Asterope* [*Cyclasterope*]
- hilgendorfii*, *Cyclasterope*
- hilgendorfii*, *Cylindroleberis* [*Cyclasterope*]
- hirpex*, *Synasterope*
- hirsutum*, *Asteropteron* [*Asteropterygion*]
- hirsutum*, *Asteropterygion*
- hirutai*, *Parasterope*
- Homasterope*
- homelix*, *Heptonema*
- hulingsi*, *Asteropteron* [*Asteropterygion*]
- hulingsi*, *Asteropterygion*
- hulingsi*, *Parasterope*
- hummelincki*, *Actinoseta*
- implumis*, *Synasterope*

- index, Synasterope*
inermis, Skogsbergiella
intesi, Pteromeniscus
jensenii, Parasterope
johanseni, Dolasterope
jonesii, Actinoseta
kalkei, Asteropella
kappa, Parasterope
kaufmani, Asteropella
keiensis, Heptonema
kliei, Cylindroleberis
kurilensis, Bathyleberis
kurilensis, Empoulsenia [Bathyleberis]
lagunicola, Parasterope
latex, Heptonema
latum, Heptonema
Leuroleberis
liguriae, Asteropterygion
liguriae, Cyclasterope [Asteropterygion]
Lobianci, Cylindroleberis [Cycloleberis]
lobiancoi, Cyclasterope [Cycloleberis]
lobiancoi, Cycloleberis
longiseta, Parasterope
longiseta, Synasterope
longungues, Parasterope
lowryi, Parasterope
lux, Parasterope
lux, Prionotoleberis
maccaini, Homasterope
mackenziei, Leuroleberis
mackenziei, Polyleberis
maclughlinae, Asteropella
Macroasteropteron
macrothrix, Skogsbergiella
maddocksa, Parasterope
maddocksa, Tetraleberis
magnum, Asteropteron [Asteropterygion]
magnum, Asteropterygion
mariae, Asterope [Cylindroleberis]
mariae, Asteropina [Cylindroleberis]
mariae, Cylindroleberis
mariae, Cypridina [Cylindroleberis]
marranyin, Cylindroleberis
mauritania, Parasterope
mckenziei, Parasterope
merx, Dromromeus
messingi, Postasterope
micra, Homasterope
Microasteropteron
microminata, Parasterope
mindax, Macroasteropteron
minuta, Asteropina [Cylindroleberis]
minuta, Cylindroleberis
monaniba, Asteropella [Asteropella monambon]
monambou, Asteropella
monambon, Asteropteron [Asteropella]
Monasterope
monothrix, Archasterope
monothrix, Bathyleberis
monothrix, Empoulsenia [Archasterope]
mortensenii, Asteropella
muelleri longiseta, Parasterope [Parasterope longiseta]
muelleri, Asteropina [Parasterope]
muelleri, Parasterope
Müller var. longiseta, Asterope [Parasterope longiseta]
Müller, Asterope [Parasterope]
mystax, Synasterope
nana, Parasterope
nodosa, Actinoseta
nodulifera, Asteropina [Cylindroleberis]
nodulifera, Cylindroleberis
nodulosum, Asteropteron [Asteropterygion]
nodulosum, Asteropterygion
norvegica, Asterope [Prionotoleberis]
norvegica, Prionotoleberis
obesa, Parasterope
oblonga, Cylindroleberis [Cylindroleberis grimaldi vicina]
oculata, Asterope [Synasterope]
oculata, Bathyleberis
oculata, Synasterope
oculitristis, Asteropteron [Asteropterygion]
oculitristis, Asteropterygion
Ohlini, Asterope [Parasterope]
ohlini, Parasterope
Omegasterope
orbicularis, Cyclasterope [Leuroleberis]
orbicularis, Cycloleberis [Leuroleberis]
orbicularis, Leuroleberis
ovalis, Cylindroleberis [Homasterope glacialis]
ovulum, Cyclasterope [Leuroleberis zealandica]
ovulum, Cycloleberis [Leuroleberis zealandica]
pacifica, Parasterope
Parasterope
parvum, Microasteropteron
pax, Asteropella
pax, Prionotoleberis
pax, Skogsbergiella
pectinata, Parasterope
pentathrix, Archasterope
pentathrix, Empoulsenia [Archasterope]
pentathrix, Philippiella [Archasterope]
persyiensis, Diasterope
peterseni, Asteropterygion
phalanx, Synasterope
Philippiella
Philippiella [Archasterope, Skogsbergiella]
physinx, Parasterope
pilosa, Diasterope
pix, Tetraleberis
plocus, Skogsbergiella
pollex, Parasterope
Polyleberis
polythrix, Archasterope
polythrix, Synasterope
Postasterope
poulsenii, Cycloleberis [Leuroleberis]
poulsenii, Leuroleberis
priacanthus, Cyclasterope [Amboleberis americana]
priacanthus, Cycloleberis [Amboleberis americana]
Prionotoleberis
procax, Diasterope
prolixa, Parasterope
pseudoquadrata, Cylindroleberis [Parasterope]
pseudoquadrata, Parasterope
psitticina, Cylindroleberis [Parasterope hulingsi]
psitticina, Cylindroleberis [Synasterope]
psitticina, Synasterope
Pteromeniscus
punctata, Asteropella
punctatum, Asteropella
quadrata, Asterope [Parasterope]
quadrata, Parasterope
quadrata, Synasterope
quadrata?, Parasterope
quatrisetosa, Synasterope
quinquesetae, Archasterope
quinquesetae, Asterope [Archasterope]
quinquesetae, Empoulsenia [Archasterope]
quinquesetae, Philippiella [Archasterope]
rangiroaensis, Cylindroleberis
rex, Prionotoleberis
romei, Asteropterygion
rotundicostata, Asteropella
rotundicostata, Cycloleberis [Asteropella]
salomani, Prionotoleberis
samoae, Asteropterygion
scammonensis, Asteropella
schmitti, Diasterope
scotti, Skogsbergiella
senex, Skogsbergiella

<i>serrata</i> , <i>Heptonema</i> [<i>Heptonema serratum</i>]	<i>Tetraleberis</i>
<i>serrata</i> , <i>Synasterope</i>	<i>thailandica</i> , <i>Cylindroleberis</i>
<i>serratum</i> , <i>Heptonema</i>	<i>thailandicum</i> , <i>Asteropteron</i> [<i>Astropterygion</i>]
<i>setiferum</i> , <i>Asteropteron</i> [<i>Astropterygion</i>]	<i>thailandicum</i> , <i>Astropterygion</i>
<i>setiferum</i> , <i>Astropterygion</i>	<i>thaliandica</i> , <i>Asteropina</i> [<i>Cylindroleberis</i>]
<i>setispara</i> , <i>Asteropina</i> [<i>Synasterope</i>]	<i>thomassini</i> , <i>Astropterygion</i>
<i>setispara</i> , <i>Synasterope</i>	<i>thrix</i> , <i>Bathyleberis</i>
<i>sharpei</i> , <i>Leuroleberis</i>	<i>thrix</i> , <i>Parasterope</i>
<i>similis</i> , <i>Cyclasterope</i> [<i>Tetraleberis</i>]	<i>toxotes</i> , <i>Bathyleberis</i>
<i>similis</i> , <i>Cycloleberis</i> [<i>Tetraleberis</i>]	<i>trebax</i> , <i>Homasterope</i>
<i>similis</i> , <i>Tetraleberis</i>	<i>tripla</i> , <i>Cyclasterope</i> [<i>Amboleberis americana</i>]
<i>skogsbergi</i> , <i>Asteropteron</i> [<i>Astropterygion</i>]	<i>tripla</i> , <i>Cycloleberis</i> [<i>Amboleberis americana</i>]
<i>skogsbergi</i> , <i>Astropterygion</i>	<i>triplex</i> , <i>Tetraleberis</i>
<i>skogsbergi</i> , <i>Philippiella</i> [<i>Skogsbergiella</i>]	<i>trithrix</i> , <i>Asteropella</i>
<i>skogsbergi</i> , <i>Skogsbergiella</i>	<i>trux</i> , <i>Xandarasterope</i>
<i>Skogsbergiella</i>	<i>unisetosa</i> , <i>Archasterope</i>
<i>slatteryi</i> , <i>Asteropella</i>	<i>unisetosa</i> , <i>Empoulsenia</i> [<i>Archasterope</i>]
<i>sohni</i> , <i>Parasterope</i>	<i>upsilon</i> , <i>Asteropteron</i> [<i>Omegasterope</i>]
<i>solox</i> , <i>Synasterope</i>	<i>upsilon</i> , <i>Omegasterope</i>
species 1, <i>Asteropella</i> [<i>Asteropella pax</i>]	<i>variabilis</i> , <i>Cylindroleberis</i>
species A, <i>Bruuniella</i> [<i>Bruuniella alpha</i>]	<i>verax</i> , <i>Archasterope</i>
species, <i>Asterope</i> [<i>Cyclasterope arthuri</i>]	<i>verrucosa</i> , <i>Asteropina</i> [<i>Cylindroleberis</i>]
<i>sphaerica</i> , <i>Cyclasterope</i> [<i>Amboleberis americana</i>]	<i>verrucosa</i> , <i>Cylindroleberis</i>
<i>sphaerica</i> , <i>Cycloleberis</i> [<i>Amboleberis americana</i>]	<i>vibex</i> , <i>Cylindroleberis</i>
<i>spinifera</i> , <i>Asterope</i> [<i>Skogsbergiella</i>]	<i>vix</i> , <i>Cylindroleberis</i>
<i>spinifera</i> , <i>Philippiella</i> [<i>Skogsbergiella</i>]	<i>weddellensis</i> , <i>Archasterope</i>
<i>spinifera</i> , <i>Skogsbergiella</i>	<i>weddellensis</i> , <i>Empoulsenia</i> [<i>Archasterope</i>]
<i>spinosum</i> , <i>Asteropteron</i> [<i>Astropterygion</i>]	<i>whatleyi</i> , <i>Parasterope</i>
c.f. <i>spinosum</i> , <i>Asteropteron</i> [<i>Astropterygion skogsbergi</i>]	<i>williamsae</i> , <i>Synasterope</i>
<i>spinosum</i> , <i>Astropterygion</i>	<i>wirraka</i> , <i>Diasterope</i>
<i>squamiger</i> , <i>Asterope</i> [<i>Cycloleberis</i>]	<i>Xandarasterope</i>
<i>squamiger</i> , <i>Cycloleberis</i>	<i>Xenoleberis</i>
<i>storthynx</i> , <i>Xandarasterope</i>	<i>yamadai</i> , <i>Bathyleberis</i> [<i>Xenoleberis</i>]
<i>styx</i> , <i>Parasterope</i>	<i>yamadai</i> , <i>Xenoleberis</i>
<i>styx</i> , <i>Prionotoleberis</i>	<i>youngi</i> , <i>Microasteropteron</i>
<i>surugaensis</i> , <i>Leuroleberis</i>	<i>zamboangae</i> , <i>Parasterope</i>
<i>Synasterope</i>	<i>zealandica</i> , <i>Azyogocypridina</i>
<i>tanzania</i> , <i>Tetraleberis</i>	<i>zealandica</i> , <i>Cyclasterope</i> [<i>Leuroleberis</i>]
<i>tenera</i> , <i>Cyclasterope</i> [<i>Leuroleberis zealandica</i>]	<i>zealandica</i> , <i>Cycloleberis</i>
<i>tenera</i> , <i>Cycloleberis</i> [<i>Leuroleberis zealandica</i>]	<i>zealandica</i> , <i>Cypridina</i> [<i>Leuroleberis</i>]
<i>tenuiseta</i> , <i>Diasterope</i>	<i>zealandica</i> , <i>Leuroleberis</i>
<i>teres</i> , <i>Cypridina</i> [<i>Parasterope muelleri</i>]	<i>zeta</i> , <i>Parasterope</i>